

CRYSTAL BASES FOR QUANTUM GENERALIZED KAC-MOODY ALGEBRAS

KYEONGHOON JEONG[◊], SEOK-JIN KANG^{*}, MASAKI KASHIWARA[†]

ABSTRACT. In this paper, we develop the crystal basis theory for quantum generalized Kac-Moody algebras. For a quantum generalized Kac-Moody algebra $U_q(\mathfrak{g})$, we first introduce the category $\mathcal{O}_{int}$ of $U_q(\mathfrak{g})$ -modules and prove its semisimplicity. Next, we define the notion of crystal bases for $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_{int}$ and for the subalgebra $U_q^-(\mathfrak{g})$. We then prove the tensor product rule and the existence theorem for crystal bases. Finally, we construct the global bases for $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_{int}$ and for the subalgebra $U_q^-(\mathfrak{g})$.

Introduction

The *quantum groups*, introduced independently by Drinfel'd and Jimbo, are certain families of Hopf algebras that are deformations of universal enveloping algebras of Kac-Moody algebras [5, 7]. More precisely, let $\mathfrak{g}$ be a symmetrizable Kac-Moody algebra and let $U(\mathfrak{g})$ be its universal enveloping algebra. Then, for each generic parameter q , we can associate a Hopf algebra $U_q(\mathfrak{g})$, called the quantum group, whose structure *tends to* that of $U(\mathfrak{g})$ as q approaches 1. In [12], Lusztig showed that the integrable highest weight modules over $U(\mathfrak{g})$ can be deformed to those over $U_q(\mathfrak{g})$ in such a way that the dimensions of weight spaces are invariant under the deformation.

In [9, 10], Kashiwara developed the *crystal basis theory* for the quantum groups associated with symmetrizable Kac-Moody algebras. (In [13], a more geometric approach was developed by Lusztig, which is called the *canonical basis theory*.) The

[◊]This research was supported by KOSEF Grant # 98-0701-01-5-L.

^{*}This research was supported by KOSEF Grant # 98-0701-01-5-L and the Young Scientist Award, Korean Academy of Science and Technology.

[†]This research is partially supported by Grant-in-Aid for Scientific Research (B1)13440006, Japan Society for the Promotion of Science.

crystal bases can be understood as bases at $q = 0$ and they are given a structure of colored oriented graphs, called the *crystal graphs*. The crystal graphs have many nice combinatorial features reflecting the internal structure of integrable modules over quantum groups. In particular, they have extremely simple behavior with respect to taking the tensor product. Thus the crystal basis theory provides us with a very powerful combinatorial tool to investigate the structure of integrable modules over quantum groups.

On the other hand, in his study of Monstrous moonshine, Borchers introduced a new class of infinite dimensional Lie algebras called the *generalized Kac-Moody algebras* [2, 3]. The structure and the representation theory of generalized Kac-Moody algebras are very similar to those of Kac-Moody algebras, and many basic facts about Kac-Moody algebras can be extended to generalized Kac-Moody algebras. But there are some differences, too. For example, generalized Kac-Moody algebras may have *imaginary simple roots* with norms ≤ 0 whose multiplicity can be greater than one. Also, they may have infinitely many simple roots.

In [8], Kang constructed the quantum groups $U_q(\mathfrak{g})$ associated with generalized Kac-Moody algebras $\mathfrak{g}$. For simplicity, we will call $U_q(\mathfrak{g})$ the *quantum generalized Kac-Moody algebras*. Moreover, he showed that, for a generic q , Verma modules and irreducible highest weight modules over $U(\mathfrak{g})$ with dominant integral highest weights can be deformed to those over $U_q(\mathfrak{g})$ in such a way that the dimensions of weight spaces are invariant under the deformation. These results were extended to generalized Kac-Moody superalgebras by Benkart, Kang and Melville [1].

In this paper, we develop the crystal basis theory for quantum generalized Kac-Moody algebras following the framework given in [10]. After reviewing some of the basic facts about quantum generalized Kac-Moody algebras and their modules, we define the category $\mathcal{O}_{int}$ consisting of integrable $U_q(\mathfrak{g})$ -modules satisfying certain conditions on their weights and show that this category is semisimple. We then define the notion of crystal bases for $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_{int}$ and prove the standard properties of crystal bases including the *tensor product rule*.

Next, we prove the existence theorem for crystal bases using Kashiwara's *grand-loop argument*, which consists of 15 interlocking inductive statements. As is the case

with Kac-Moody algebras, the fundamental properties of crystal bases for $U_q^-(\mathfrak{g})$ play a crucial role in this argument.

Finally, we *globalize* the theory of crystal bases. That is, we show that any irreducible highest weight $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ has a unique *global basis*. The existence of global bases is proved using the basic properties of *balanced triple* and the triviality of vector bundles over $\mathbf{P}^1$.

Acknowledgments. We would like to express our sincere gratitude to Olivier Schiffmann for many valuable discussions.

1. Quantum Generalized Kac-Moody Algebras

Let I be a finite or countably infinite index set. A real square matrix $A = (a_{ij})_{i,j \in I}$ is called a *Borcherds-Cartan matrix* if it satisfies:

- (i) $a_{ii} = 2$ or $a_{ii} \leq 0$ for all $i \in I$,
- (ii) $a_{ij} \leq 0$ if $i \neq j$,
- (iii) $a_{ij} \in \mathbf{Z}$ if $a_{ii} = 2$,
- (iv) $a_{ij} = 0$ if and only if $a_{ji} = 0$.

We say that an index i is *real* if $a_{ii} = 2$ and *imaginary* if $a_{ii} \leq 0$. We denote by $I^{\text{re}} := \{i \in I; a_{ii} = 2\}$, $I^{\text{im}} := \{i \in I; a_{ii} \leq 0\}$.

In this paper, we assume that all the entries of A are integers and the diagonal entries are even. Furthermore, we assume that A is *symmetrizable*; that is, there is a diagonal matrix $D = \text{diag}(s_i \in \mathbf{Z}_{>0} \mid i \in I)$ such that DA is symmetric.

Let $P^\vee$ be a free abelian group and let $\Pi^\vee = \{h_i\}_{i \in I}$ be a family of elements in $P^\vee$ called the set of *simple coroots*, and set $\mathfrak{h} = \mathbf{Q} \otimes_{\mathbf{Z}} P^\vee$. The free abelian group $P^\vee$ is called the *co-weight lattice* and the $\mathbf{Q}$ -vector space $\mathfrak{h}$ is called the *Cartan subalgebra*. The *weight lattice* is defined to be

$$P := \{\lambda \in \mathfrak{h}^*; \lambda(P^\vee) \subset \mathbf{Z}\}.$$

We assume that P contains a family of elements $\Pi = \{\alpha_i\}_{i \in I}$ called the set of *simple roots* satisfying

$$\alpha_i(h_j) = a_{ji} \quad \text{for } i, j \in I.$$

Definition 1.1. We call the quintuple $(A, P^\vee, P, \Pi^\vee, \Pi)$ a *Borcherds-Cartan datum* associated with A .

In this paper, we assume that there is a non-degenerate symmetric bilinear form $(\mid)$ on $\mathfrak{h}^*$ satisfying

$$(\alpha_i \mid \lambda) = s_i \lambda(h_i) \quad \text{for every } \lambda \in \mathfrak{h}^*,$$

and therefore we have

$$(\alpha_i \mid \alpha_j) = s_i a_{ij} \quad \text{for every } i, j \in I.$$

We also assume that there exists $\Lambda_i \in P$ such that $\Lambda_i(h_j) = \delta_{ij}$ for $i, j \in I$. Such a Λ_i is called a *fundamental weight*. We assume further that $\{\alpha_i\}_{i \in I}$ is linearly independent.

For a symmetrizable Borcherds-Cartan matrix, there exists such a Borcherds-Cartan datum. Take

$$P^\vee = \left(\bigoplus_{i \in I} \mathbf{Z} h_i \right) \oplus \left(\bigoplus_{i \in I} \mathbf{Z} d_i \right)$$

and define α_i and Λ_i by

$$\begin{aligned} \alpha_i(h_j) &= a_{ji} \text{ and } \alpha_i(d_j) = \delta_{ij}, \\ \Lambda_i(h_j) &= \delta_{ij} \text{ and } \Lambda_i(d_j) = 0. \end{aligned}$$

Let $(\mid)$ be the bilinear form on $(\bigoplus_i (\mathbf{Q}\alpha_i \oplus \mathbf{Q}\Lambda_i)) \times \mathfrak{h}^*$ defined by

$$(\alpha_i \mid \lambda) = s_i \lambda(h_i), \quad (\Lambda_i \mid \lambda) = s_i \lambda(d_i).$$

Since it is symmetric on $(\bigoplus_i (\mathbf{Q}\alpha_i \oplus \mathbf{Q}\Lambda_i)) \times (\bigoplus_i (\mathbf{Q}\alpha_i \oplus \mathbf{Q}\Lambda_i))$, we can extend this to a symmetric form on $\mathfrak{h}^*$. Then such a symmetric form is non-degenerate.

We denote by P^+ the set $\{\lambda \in P; \lambda(h_i) \geq 0 \text{ for every } i \in I\}$ of *dominant integral weights*. The free abelian group $Q := \bigoplus_{i \in I} \mathbf{Z}\alpha_i$ is called the *root lattice*. We set $Q_+ = \sum_{i \in I} \mathbf{Z}_{\geq 0} \alpha_i$ and $Q_- = -Q_+$. For $\alpha \in Q_+$, we can write $\alpha = \sum_{k=1}^r \alpha_{i_k}$ for $i_1, \dots, i_r \in I$. We set $|\alpha| = r$ and call it the *height* of α .

For an indeterminate q , set $q_i = q^{s_i}$ and define

$$(1.1) \quad [n]_i = \frac{q_i^n - q_i^{-n}}{q_i - q_i^{-1}}, \quad [n]_i! = \prod_{k=1}^n [k]_i, \quad \left[\begin{matrix} m \\ n \end{matrix} \right]_i = \frac{[m]_i!}{[m-n]_i! [n]_i!}.$$

If $a_{ii} < 0$, set $c_i = -\frac{1}{2}a_{ii} \in \mathbf{Z}_{>0}$, and define

$$(1.2) \quad \{n\}_i = \frac{q_i^{c_i n} - q_i^{-c_i n}}{q_i^{c_i} - q_i^{-c_i}}, \quad \{n\}_i! = \prod_{k=1}^n \{k\}_i, \quad \left\{ \begin{matrix} m \\ n \end{matrix} \right\}_i = \frac{\{m\}_i!}{\{m-n\}_i! \{n\}_i!}.$$

For convenience, if $a_{ii} = 0$, we will also use the notation $c_i = 0$ and

$$(1.3) \quad \{n\}_i = n, \quad \{n\}_i! = n!, \quad \left\{ \begin{matrix} m \\ n \end{matrix} \right\}_i = \binom{m}{n}.$$

Definition 1.2. The *quantum generalized Kac-Moody algebra* $U_q(\mathfrak{g})$ associated with a Borcherds-Cartan datum $(A, P^\vee, P, \Pi^\vee, \Pi)$ is the associative algebra over $\mathbf{Q}(q)$ with the unit 1 generated by the symbols e_i, f_i ($i \in I$) and q^h ($h \in P^\vee$) subject to the following defining relations:

$$(1.4) \quad \begin{aligned} q^0 &= 1, \quad q^h q^{h'} = q^{h+h'} \quad (h, h' \in P^\vee), \\ q^h e_i q^{-h} &= q^{\alpha_i(h)} e_i, \quad q^h f_i q^{-h} = q^{-\alpha_i(h)} f_i, \\ e_i f_j - f_j e_i &= \delta_{ij} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}, \quad \text{where } K_i = q^{s_i h_i}, \\ \sum_{r=0}^{1-a_{ij}} (-1)^r \left[\begin{matrix} 1-a_{ij} \\ r \end{matrix} \right]_i e_i^{1-a_{ij}-r} e_j e_i^r &= 0 \quad \text{if } a_{ii} = 2, i \neq j, \\ \sum_{r=0}^{1-a_{ij}} (-1)^r \left[\begin{matrix} 1-a_{ij} \\ r \end{matrix} \right]_i f_i^{1-a_{ij}-r} f_j f_i^r &= 0 \quad \text{if } a_{ii} = 2, i \neq j, \\ e_i e_j - e_j e_i &= 0, \quad f_i f_j - f_j f_i = 0 \quad \text{if } a_{ij} = 0. \end{aligned}$$

From now on, we will use the notation

$$(1.5) \quad \begin{aligned} e_i^{(n)} &= \frac{e_i^n}{[n]_i!}, \quad f_i^{(n)} = \frac{f_i^n}{[n]_i!} \quad \text{if } a_{ii} = 2, \\ e_i^{(n)} &= e_i^n, \quad f_i^{(n)} = f_i^n \quad \text{if } a_{ii} \leq 0. \end{aligned}$$

We understand that $e_i^{(n)} = f_i^{(n)} = 1$ for $n = 0$ and $e_i^{(n)} = f_i^{(n)} = 0$ for $n < 0$. For $\alpha = \sum k_i \alpha_i \in Q$, we will also use the notation

$$K_\alpha = \prod K_i^{k_i}.$$

The quantum generalized Kac-Moody algebra $U_q(\mathfrak{g})$ has a *Hopf algebra* structure with the comultiplication Δ , the counit ε , and the antipode S defined by

$$(1.6) \quad \begin{aligned} \Delta(q^h) &= q^h \otimes q^h, \\ \Delta(e_i) &= e_i \otimes K_i^{-1} + 1 \otimes e_i, \quad \Delta(f_i) = f_i \otimes 1 + K_i \otimes f_i, \\ \varepsilon(q^h) &= 1, \quad \varepsilon(e_i) = \varepsilon(f_i) = 0, \\ S(q^h) &= q^{-h}, \quad S(e_i) = -e_i K_i, \quad S(f_i) = -K_i^{-1} f_i \end{aligned}$$

for $h \in P^\vee$, $i \in I$ (see, for example, [1, 8]).

Let $U_q^+(\mathfrak{g})$ (resp. $U_q^-(\mathfrak{g})$) be the subalgebra of $U_q(\mathfrak{g})$ generated by the elements e_i (resp. f_i) for $i \in I$, and let $U_q^0(\mathfrak{g})$ be the subalgebra of $U_q(\mathfrak{g})$ generated by q^h ($h \in P^\vee$). Then we have the *triangular decomposition* ([1, 8]):

$$(1.7) \quad U_q(\mathfrak{g}) \cong U_q^-(\mathfrak{g}) \otimes U_q^0(\mathfrak{g}) \otimes U_q^+(\mathfrak{g}).$$

Remark 1.3. In [8], Kang considered the generalized Kac-Moody algebras associated with Borcherds-Cartan matrices of *charge* $\mathbf{m} = (m_i \in \mathbf{Z}_{>0} \mid i \in I, \ m_i = 1 \text{ for } i \in I^{\text{re}})$. The charge m_i is the multiplicity of the simple root corresponding to $i \in I$. For example, the *Monster Lie algebra*, which played an important role in Borcherds' proof of Monstrous moonshine, is a generalized Kac-Moody algebra associated with the Borcherds-Cartan matrix $A = (-(i+j))_{i,j \in I}$ of charge $\mathbf{m} = (c(i) \mid i \in I)$, where $I = \{-1\} \cup \{1, 2, \dots\}$ and $c(i)$ are the coefficients of the elliptic modular function

$$J(q) = j(q) - 744 = \sum_{n=-1}^{\infty} c(n)q^n = q^{-1} + 196884q + 21493760q^2 + \dots$$

In this paper, we assume that $m_i = 1$ for all $i \in I$. However, we do not lose generality by this assumption. Indeed, if we take Borcherds-Cartan matrices with some of the rows and columns identical, then the generalized Kac-Moody algebras with charge introduced in [8] can be recovered from the ones in this paper by identifying the h_i 's and d_i 's (hence α_i 's) corresponding to these identical rows and columns.

2. Representation Theory

A $U_q(\mathfrak{g})$ -module V is called a *weight module* if it admits a *weight space decomposition* $V = \bigoplus_{\mu \in P} V_\mu$, where $V_\mu := \{v \in V; q^h v = q^{\mu(h)} v \text{ for every } h \in P^\vee\}$. We call $\text{wt}(V) := \{\mu \in P; V_\mu \neq 0\}$ the set of *weights* of V .

Let $\mathcal{O}$ be the category consisting of weight modules V with finite-dimensional weight spaces such that $\text{wt}(V) \subset \bigcup_{j=1}^s (\lambda_j - Q_+)$ for finitely many $\lambda_1, \dots, \lambda_s \in P$.

One of the most interesting examples of $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}$ is the class of highest weight modules defined below. A weight module V over $U_q(\mathfrak{g})$ is called a *highest weight module with highest weight* $\lambda \in P$ if there exists a *non-zero* vector $v \in V$ (called a *highest weight vector*) such that

- (i) $V = U_q(\mathfrak{g})v$,
- (ii) $v \in V_\lambda$,
- (iii) $e_i v = 0$ for every $i \in I$.

Let $J(\lambda)$ denote the left ideal of $U_q(\mathfrak{g})$ generated by $e_i, q^h - q^{\lambda(h)}$ ($i \in I, h \in P^\vee$), and set $M(\lambda) = U_q(\mathfrak{g})/J(\lambda)$. Then, via left multiplication, $M(\lambda)$ becomes a $U_q(\mathfrak{g})$ -module called the *Verma module*. The basic properties of Verma modules are summarized in the following proposition.

Proposition 2.1 ([1, 8]).

- (a) $M(\lambda)$ is a highest weight $U_q(\mathfrak{g})$ -module with highest weight λ and highest weight vector $v_\lambda := 1 + J(\lambda)$.
- (b) $M(\lambda)$ is a free $U_q^-(\mathfrak{g})$ -module generated by v_λ .
- (c) Every highest weight $U_q(\mathfrak{g})$ -module with highest weight λ is a quotient of $M(\lambda)$.
- (d) $M(\lambda)$ contains a unique maximal submodule $R(\lambda)$.

The irreducible quotient $V(\lambda) = M(\lambda)/R(\lambda)$ is an *irreducible highest weight* $U_q(\mathfrak{g})$ -module with highest weight λ .

In the rest of this section, we focus on the structure of the irreducible highest weight $U_q(\mathfrak{g})$ -module $V(\lambda)$ with a dominant integral highest weight $\lambda \in P^+$.

We first recall:

Proposition 2.2 ([1, 8]). *Let $\lambda \in P^+$ be a dominant integral weight.*

(a) *The highest weight vector v_λ of $V(\lambda)$ satisfies the following relations*

$$(2.1) \quad \begin{cases} f_i^{\lambda(h_i)+1} v_\lambda = 0 & \text{if } a_{ii} = 2, \\ f_i v_\lambda = 0 & \text{if } \lambda(h_i) = 0. \end{cases}$$

(b) *Conversely, let V be a highest weight $U_q(\mathfrak{g})$ -module with a highest weight $\lambda \in P^+$ and a highest weight vector v . If v satisfies the relations in (2.1), then V is isomorphic to $V(\lambda)$.*

Consider the anti-involution of $U_q(\mathfrak{g})$ defined by

$$(2.2) \quad q^h \mapsto q^h, \quad e_i \mapsto q_i f_i K_i^{-1}, \quad f_i \mapsto q_i^{-1} K_i e_i \quad (i \in I).$$

By standard arguments, we see that there exists a unique symmetric bilinear form $(\ , \)$ on $V(\lambda)$ with $\lambda \in P^+$ satisfying

$$(2.3) \quad \begin{aligned} (v_\lambda, v_\lambda) &= 1, \quad (q^h u, v) = (u, q^h v), \\ (e_i u, v) &= (u, q_i f_i K_i^{-1} v), \quad (f_i u, v) = (u, q_i^{-1} K_i e_i v) \end{aligned}$$

for $i \in I$, $h \in P^\vee$, $u, v \in V(\lambda)$.

Note that $(V(\lambda)_\mu, V(\lambda)_\tau) = 0$ unless $\mu = \tau$. Moreover, we have:

Lemma 2.3. *The symmetric bilinear form $(\ , \)$ on $V(\lambda)$ defined by (2.3) is non-degenerate.*

Proof. The kernel $\{u \in V(\lambda) ; (u, V(\lambda)) = 0\}$ is a $U_q(\mathfrak{g})$ -submodule not containing v_λ . Since $V(\lambda)$ is irreducible, it must vanish. $\square$

The weights of the irreducible highest weight $U_q(\mathfrak{g})$ -module $V(\lambda)$ satisfy the following properties.

Proposition 2.4. *Let μ be a weight of $V(\lambda)$ with $\lambda \in P^+$, and let $i \in I^{\text{im}}$.*

- (a) $\mu(h_i) \in \mathbf{Z}_{\geq 0}$.
- (b) If $\mu(h_i) = 0$, then $V(\lambda)_{\mu - \alpha_i} = 0$.
- (c) If $\mu(h_i) = 0$, then $f_i(V(\lambda)_\mu) = 0$.
- (d) If $\mu(h_i) \leq 2c_i$, then $e_i(V(\lambda)_\mu) = 0$.

Proof. Set $\mu = \lambda - \alpha$ with $\alpha \in Q_+$. Write $\alpha = \sum_{\nu=1}^r \alpha_{i_\nu}$. Since $\alpha_{i_\nu}(h_i) \leq 0$, we have $\alpha(h_i) \leq 0$, which yields $\mu(h_i) = \lambda(h_i) - \alpha(h_i) \geq \lambda(h_i) \geq 0$. Note that $\mu(h_i) = 0$ implies that $\lambda(h_i) = \alpha_{i_\nu}(h_i) = 0$ ($\nu = 1, \dots, r$). Thus f_i commutes with all the f_{i_ν} 's, and kills v_λ . Therefore we obtain $V(\lambda)_{\mu-\alpha_i} = 0$.

The properties (c) and (d) easily follow from (a) and (b). $\square$

Fix $i \in I$, and let U_i be the subalgebra of $U_q(\mathfrak{g})$ generated by e_i , f_i and $U_q^0(\mathfrak{g})$. Then the algebra U_i can be viewed as a rank 1 quantum generalized Kac-Moody algebra associated with the Borcherds-Cartan matrix $A = (a_{ii})$. For example, if $a_{ii} = 0$, then U_i is isomorphic to the *quantum Heisenberg algebra*.

Let $V = V(\lambda)$ be the irreducible highest weight U_i -module with a dominant integral highest weight λ . If $a_{ii} = 2$, it is well-known that $\dim V = \lambda(h_i) + 1$. If $a_{ii} \leq 0$, then $\dim V$ is given in the following lemma, which easily follows from Proposition 2.2.

Lemma 2.5. *Suppose $a_{ii} \leq 0$ and let $V(\lambda)$ be the irreducible highest weight U_i -module with a dominant integral highest weight λ . Then we have $V(\lambda) = \mathbf{Q}(q)v_\lambda$ if $\lambda(h_i) = 0$, and $\{f_i^n v_\lambda\}_{n \geq 0}$ is a basis of $V(\lambda)$ if $\lambda(h_i) > 0$.*

Observe that the U_i -module structure on $V(\lambda)$ is given by

$$(2.4) \quad \begin{aligned} K_i f_i^n v_\lambda &= q_i^{\lambda(h_i) + 2nc_i} f_i^n v_\lambda, \\ f_i f_i^n v &= f_i^{n+1} v, \\ e_i f_i^n v_\lambda &= \{n\}_i [\lambda(h_i) + c_i(n-1)] f_i^{n-1} v_\lambda. \end{aligned}$$

3. Category $\mathcal{O}_{int}$

Definition 3.1. The category $\mathcal{O}_{int}$ consists of $U_q(\mathfrak{g})$ -modules M satisfying the following properties:

- (i) $M = \bigoplus_{\mu \in P} M_\mu$ with $\dim M_\mu < \infty$ for every $\mu \in P$, where

$$M_\mu := \{v \in M; q^h v = q^{\mu(h)} v \text{ for every } h \in P^\vee\},$$

(ii) there exist a finite number of weights $\lambda_1, \dots, \lambda_s \in P$ such that

$$\text{wt}(M) := \{\mu \in P; M_\mu \neq 0\} \subset \bigcup_{j=1}^s (\lambda_j - Q_+),$$

(iii) if $a_{ii} = 2$, then the action of f_i on M is locally nilpotent,

(iv) if $a_{ii} \leq 0$, then $\mu(h_i) \in \mathbf{Z}_{\geq 0}$ for every $\mu \in \text{wt}(M)$,

(v) if $a_{ii} \leq 0$ and $\mu(h_i) = 0$, then $f_i M_\mu = 0$.

(vi) if $a_{ii} \leq 0$ and $\mu(h_i) = 2c_i$, then $e_i M_\mu = 0$.

Remark 3.2.

- (i) Note that if $a_{ii} \leq 0$ and $\mu(h_i) \leq 2c_i$ then $e_i M_\mu = 0$. Indeed, if $\mu(h_i) < 2c_i$, then $(\mu + \alpha_i)(h_i) < 0$ and $\mu + \alpha_i$ is not a weight of M by Definition 3.1 (iv).
- (ii) By (ii), the action of e_i on a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ is locally nilpotent.
- (iii) Note that a submodule or a quotient module of a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ also belongs to the category $\mathcal{O}_{int}$. Furthermore, it is straightforward to verify that the category $\mathcal{O}_{int}$ is closed under taking a finite number of direct sums and tensor products.

Proposition 3.3. *Let $V(\lambda)$ be the irreducible highest weight $U_q(\mathfrak{g})$ -module with highest weight $\lambda \in P$. Then $V(\lambda)$ belongs to the category $\mathcal{O}_{int}$ if and only if $\lambda \in P^+$.*

Proof. As we have seen in Proposition 2.4, $V(\lambda)$ belongs to the category $\mathcal{O}_{int}$ for $\lambda \in P^+$.

Conversely, assume that $V(\lambda)$ lies in the category $\mathcal{O}_{int}$. If $a_{ii} = 2$, since f_i is locally nilpotent on $V(\lambda)$, there exists a non-negative integer N_i such that $f_i^{N_i} v_\lambda \neq 0$ and $f_i^{N_i+1} v_\lambda = 0$. Hence we have

$$0 = e_i f_i^{N_i+1} v_\lambda = [N_i + 1]_i [\lambda(h_i) - N_i]_i f_i^{N_i} v_\lambda,$$

which implies $\lambda(h_i) = N_i \in \mathbf{Z}_{\geq 0}$.

If $a_{ii} \leq 0$, we have $\lambda(h_i) \in \mathbf{Z}_{\geq 0}$ by Definition 3.1 (iv). □

Proposition 3.4.

- (a) If V is a highest weight $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ with highest weight $\lambda \in P$, then $\lambda \in P^+$ and $V \simeq V(\lambda)$.
- (b) Every irreducible $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ is isomorphic to some $V(\lambda)$ with $\lambda \in P^+$.

Proof. (a) Let v be a highest weight vector of V with weight λ . If $a_{ii} = 2$, by the proof of Proposition 3.3, we have $\lambda(h_i) \in \mathbf{Z}_{\geq 0}$ and $f_i^{\lambda(h_i)+1}v = 0$.

If $a_{ii} \leq 0$, then by the definition of $\mathcal{O}_{int}$, we have $\lambda(h_i) \in \mathbf{Z}_{\geq 0}$ and $\lambda(h_i) = 0$ implies $f_i v = 0$. Hence $\lambda \in P^+$, and Proposition 2.2 (b) implies that $V \simeq V(\lambda)$.

(b) Let V be an irreducible $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$. Since $\text{wt}(V) \subset \bigcup_{j=1}^s (\lambda_j - Q_+)$ for some $\lambda_j \in P$ ($j = 1, \dots, s$), there exists a non-zero *maximal vector* $v \in V_\lambda$ for some $\lambda \in \text{wt}(V) \subset P$ such that $e_i v = 0$ for every $i \in I$. Then $V = U_q(\mathfrak{g})v$ is isomorphic to $V(\lambda)$ by (a). $\square$

We will now prove that every $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ is semisimple. Let φ be the anti-involution of $U_q(\mathfrak{g})$ given by

$$(3.1) \quad e_i \mapsto f_i, \quad f_i \mapsto e_i, \quad q^h \mapsto q^h.$$

Let $M = \bigoplus_{\mu \in P} M_\mu$ be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$. We define its *finite dual* to be the vector space

$$(3.2) \quad M^* := \bigoplus_{\mu \in P} M_\mu^*, \quad \text{where } M_\mu^* := \text{Hom}_{\mathbf{Q}(q)}(M_\mu, \mathbf{Q}(q)).$$

Using the anti-involution φ of $U_q(\mathfrak{g})$, we define a $U_q(\mathfrak{g})$ -module structure on M^* by

$$(3.3) \quad \langle x \cdot \phi, v \rangle = \langle \phi, \varphi(x) \cdot v \rangle \quad \text{for } x \in U_q(\mathfrak{g}), \phi \in M^*, v \in M.$$

The following lemma is an immediate consequence of the definitions.

Lemma 3.5. *Let $M = \bigoplus_{\mu \in P} M_\mu$ be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$.*

- (a) *There exists a canonical isomorphism $(M^*)^* \cong M$ as $U_q(\mathfrak{g})$ -modules.*
- (b) *The space M_μ^* is the weight space of M^* with weight μ .*
- (c) *We have*

$$\text{wt}(M^*) = \text{wt}(M).$$

(d) M^* also belongs to the category $\mathcal{O}_{int}$.

Proof. Since the other statements are evident, we only show that M^* satisfies (v) and (vi) in Definition 3.1.

(v) For $i \in I^{\text{im}}$ and $\mu \in P$ with $\mu(h_i) = 0$, we have

$$\langle f_i M_\mu^*, M \rangle = \langle M_\mu^*, e_i M_{\mu - \alpha_i} \rangle = 0,$$

where the last equality follows from $\langle h_i, \mu - \alpha_i \rangle = 2c_i$.

(vi) For $i \in I^{\text{im}}$ and $\mu \in P$ with $\mu(h_i) = 2c_i$, we have

$$\langle e_i M_\mu^*, M \rangle = \langle M_\mu^*, f_i M_{\mu + \alpha_i} \rangle = 0,$$

where the last equality follows from $\langle h_i, \mu + \alpha_i \rangle = 0$. □

Let M be a non-zero $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ and choose a *maximal weight* $\lambda \in \text{wt}(M)$ with the property that $\lambda + \alpha_i$ is *not* a weight of M for any $i \in I$. Fix a non-zero vector $v_\lambda \in M_\lambda$ and set $V = U_q(\mathfrak{g})v_\lambda$. Then by Proposition 3.4, we have $\lambda \in P^+$ and V is isomorphic to $V(\lambda)$.

Let us take $\phi_\lambda \in M_\lambda^*$ satisfying $\phi_\lambda(v_\lambda) = 1$, and set $W = U_q(\mathfrak{g})\phi_\lambda$. Then, again by Proposition 3.4, W is isomorphic to $V(\lambda)$.

Lemma 3.6. *Let M be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ and let V be the submodule of M generated by a maximal vector v_λ of weight λ . Then we have*

$$M \simeq V \oplus (M/V).$$

Proof. We will show that the short exact sequence

$$0 \longrightarrow V \longrightarrow M \longrightarrow M/V \longrightarrow 0$$

splits.

Take the dual of the inclusion $W \hookrightarrow M^*$ to get a map $(M^*)^* \longrightarrow W^*$. Since $M \cong (M^*)^*$, we obtain a map $\eta: M \xrightarrow{\sim} (M^*)^* \longrightarrow W^*$, which yields

$$\psi: V \hookrightarrow M \xrightarrow{\eta} W^*.$$

It is easy to check that the image of v_λ under ψ is non-zero. Since $V \simeq V(\lambda)$ and $W^* \simeq V(\lambda)^* \simeq V(\lambda)$, ψ is an isomorphism by Schur's Lemma. Hence the above short exact sequence splits, and we have $M \simeq V \oplus (M/V)$. $\square$

Using this lemma, we will prove the complete reducibility theorem for the category $\mathcal{O}_{int}$.

Theorem 3.7. *Every $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ is isomorphic to a direct sum of irreducible highest weight modules $V(\lambda)$ with $\lambda \in P^+$.*

Proof. By Proposition 3.4 (b), it is enough to show that any $U_q(\mathfrak{g})$ -module M in the category $\mathcal{O}_{int}$ is semisimple.

Let $U^{\geq 0}$ denote the subalgebra of $U_q(\mathfrak{g})$ generated by the elements q^h ($h \in P^\vee$) and e_i ($i \in I$). We shall first show that, if M is generated as a $U_q(\mathfrak{g})$ -module by a finite-dimensional $U^{\geq 0}$ -submodule F , then M is semisimple. We shall argue by the induction on the dimension of F . If $F \neq 0$, let us choose a maximal weight vector v_λ of weight $\lambda \in P^+$ in F , and set $V = U_q(\mathfrak{g})v_\lambda \simeq V(\lambda)$. By Lemma 3.6, we have

$$M \simeq V \oplus (M/V).$$

Since $M/V = U_q(\mathfrak{g})(F/(F \cap V))$ and $\dim(F/(F \cap V)) < \dim F$, the induction hypothesis implies that M/V is semisimple. Hence M is semisimple.

Now let M be an arbitrary $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$. Then, for any $v \in M$, $U^{\geq 0}v$ is finite-dimensional, and hence $U_q(\mathfrak{g})v$ is semisimple. Since M is a sum of semisimple $U_q(\mathfrak{g})$ -modules $U_q(\mathfrak{g})v$, a standard argument for semisimplicity ([4, Proposition 3.12]) implies that M is semisimple. $\square$

Corollary 3.8. *Let M be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$. Then for any $i \in I$, M satisfies the following properties:*

- (a) *If $a_{ii} = 2$, then M is a direct sum of finite-dimensional irreducible U_i -submodules.*
- (b) *If $a_{ii} \leq 0$, then M is a direct sum of 1-dimensional or infinite-dimensional irreducible highest weight U_i -submodules.*

Proof. Our assertions immediately follow from Lemma 2.5 and Theorem 3.7. $\square$

4. Crystal Bases

In this section, we will develop the *crystal basis theory* for $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_{int}$. The following fundamental lemma easily follows from Corollary 3.8.

Lemma 4.1. *Let M be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$. Then for any $i \in I$, we have:*

- (a) $M = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} f_i^{(n)}(\text{Ker } e_i)$.
- (b) For any $\lambda \in P$ and $n > 0$, the map

$$(4.1) \quad f_i^{(n)} : \text{Ker } e_i \cap M_\lambda \rightarrow M_{\lambda - n\alpha_i}$$

is either a monomorphism or zero. More precisely, the morphism (4.1) is injective if $i \in I^{\text{re}}$ and $\lambda(h_i) \geq n$ or if $i \in I^{\text{im}}$ and $\lambda(h_i) > 0$. Otherwise, it vanishes.

Hence for any weight vector $u \in M_\lambda$ there exists a unique family $\{u_n\}_{n \in \mathbb{Z}_{\geq 0}}$ of elements of M such that

$$(4.2) \quad \begin{aligned} & \text{(a) } u_n \in \text{Ker } e_i \cap M_{\lambda + n\alpha_i}, \\ & \text{(b) } u = \sum_{n \geq 0} f_i^{(n)} u_n, \\ & \text{(c) if } i \in I^{\text{re}} \text{ and } \langle h_i, \lambda + n\alpha_i \rangle < n, \text{ then } u_n = 0, \\ & \text{(d) if } i \in I^{\text{im}}, n > 0 \text{ and } \langle h_i, \lambda + n\alpha_i \rangle = 0, \text{ then } u_n = 0. \end{aligned}$$

Note that $u_n = 0$ for $n \gg 0$, because $\lambda + n\alpha_i$ is not a weight of M for $n \gg 0$. We call the expression (b) in (4.2) the *i-string decomposition* of u .

Remark 4.2.

- (i) The conditions (c) and (d) in (4.2) are equivalent to saying that $f_i^{(n)} u_n = 0$ implies $u_n = 0$.
- (ii) If $\sum_{n \geq 0} f_i^{(n)} u_n$ is an *i-string decomposition*, then $\sum_{n \geq 0} f_i^{(n)} u_{n+1}$ is an *i-string decomposition*. However, $\sum_{n \geq 1} f_i^{(n)} u_{n-1}$ is not necessarily an *i-string decomposition*.

Definition 4.3. For $i \in I$ and $u \in M$, let $u = \sum_{n \geq 0} f_i^{(n)} u_n$ be the i -string decomposition of u . Then the *Kashiwara operators* $\tilde{e}_i$ and $\tilde{f}_i$ on M are defined by

$$(4.3) \quad \tilde{e}_i u = \sum_{n \geq 1} f_i^{(n-1)} u_n, \quad \tilde{f}_i u = \sum_{n \geq 0} f_i^{(n+1)} u_n.$$

By Lemma 4.1, if $u = \sum_{n \geq 0} f_i^{(n)} u_n$ with $u \in M_\lambda$ and $u_n \in \text{Ker } e_i \cap M_{\lambda+n\alpha_i}$, then we have

$$(4.4) \quad \tilde{e}_i u = \sum_{\substack{n \geq 1, \\ f_i^{(n)} u_n \neq 0}} f_i^{(n-1)} u_n \quad \text{and} \quad \tilde{f}_i u = \sum_{n \geq 0} f_i^{(n+1)} u_n.$$

Note that if $a_{ii} \leq 0$, then $\tilde{f}_i = f_i$, but $\tilde{e}_i \neq e_i$ (see Lemma 4.5 below and the remark after (4.6)). Note that we have

$$(4.5) \quad \tilde{e}_i \circ \tilde{f}_i |_{M_\lambda} = \text{id}_{M_\lambda} \text{ if } a_{ii} \leq 0 \text{ and } \lambda(h_i) > 0.$$

We now define the notion of *crystal bases* for $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_{int}$. Let $\mathbf{A}_0$ be the localization of the polynomial ring $\mathbf{Q}[q]$ at the prime ideal $\mathbf{Q}[q]q$:

$$\mathbf{A}_0 = \{f/g; f, g \in \mathbf{Q}[q], g(0) \neq 0\}.$$

Thus the local ring $\mathbf{A}_0$ is a principal ideal domain with $\mathbf{Q}(q)$ as its field of quotients.

Definition 4.4. Let M be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$. A *crystal lattice* of M is a free $\mathbf{A}_0$ -submodule L of M satisfying the following conditions:

- (i) L generates M as a vector space over $\mathbf{Q}(q)$,
- (ii) $L = \bigoplus_{\lambda \in P} L_\lambda$, where $L_\lambda := L \cap M_\lambda$,
- (iii) $\tilde{e}_i L \subset L$ and $\tilde{f}_i L \subset L$ for every $i \in I$.

The map $\mathbf{A}_0 \ni f \mapsto f(0) \in \mathbf{Q}$ induces an isomorphism

$$\mathbf{A}_0/q\mathbf{A}_0 \xrightarrow{\sim} \mathbf{Q},$$

and we have

$$\mathbf{Q} \otimes_{\mathbf{A}_0} L \xrightarrow{\sim} L/qL.$$

The passage from L to the quotient L/qL is referred to as taking the *crystal limit*. Since the Kashiwara operators $\tilde{e}_i$ and $\tilde{f}_i$ preserve the crystal lattice L , they also define operators on L/qL , which we will denote by the same symbols.

Lemma 4.5. *Let L be a free $\mathbf{A}_0$ -submodule L of $M \in \mathcal{O}_{int}$ satisfying the conditions (i) and (ii) in Definition 4.4.*

- (a) *L is a crystal lattice if and only if $L = \bigoplus_{n \geq 0} f_i^{(n)}(\text{Ker } e_i \cap L)$ for every $i \in I$.*
- (b) *Let E_i and F_i be operators on M such that*

$$E_i f_i^{(n)} v = a_{i,n,\lambda} f_i^{(n-1)} v, \quad F_i f_i^{(n)} v = b_{i,n,\lambda} f_i^{(n+1)} v$$

for $v \in \text{Ker } e_i \cap M_\lambda$ with $f_i^{(n)} v \neq 0$. Here, $a_{i,n,\lambda}$ and $b_{i,n,\lambda}$ are invertible elements of $\mathbf{A}_0$.

Then L is a crystal lattice if and only if $E_i L \subset L$ and $F_i L \subset L$. Moreover, if we define the operators R_i and S_i by

$$R_i f_i^{(n)} v = a_{i,n,\lambda}(0) f_i^{(n)} v, \quad S_i f_i^{(n)} v = b_{i,n,\lambda}(0) f_i^{(n)} v,$$

then $R_i L \subset L$, $S_i L \subset L$ and the induced action of E_i (resp. F_i) on L/qL coincides with that of $\tilde{e}_i \circ R_i$ (resp. $\tilde{f}_i \circ S_i$). In particular, we have $E_i^{-1}(L) = \tilde{e}_i^{-1}(L)$.

Proof. Since the other statements are obvious, let us show that if $E_i L \subset L$ and $F_i L \subset L$, then $L = \bigoplus_{n \geq 0} f_i^{(n)}(\text{Ker } e_i \cap L)$. The condition $F_i L \subset L$ implies that $L \supset \bigoplus_{n \geq 0} f_i^{(n)}(\text{Ker } e_i \cap L)$. In order to show the inverse inclusion, let $\sum_{n=0}^N f_i^{(n)} u_n$ be the i -string decomposition of $u \in L_\lambda$. Let us show that every u_n belongs to L by the induction of N . Since we have

$$E_i u = \sum_{n=1}^N a_{i,n,\lambda+n\alpha_i} f_i^{(n-1)} u_n \in L,$$

the induction hypothesis implies $u_n \in L$ for $n \geq 1$. Hence $u_0 = u - \sum_{n \geq 1} f_i^{(n)} u_n$ belongs to L . $\square$

Note that for $u \in \text{Ker } e_i \cap M_\lambda$, we have

$$(4.6) \quad q_i^{-1} K_i e_i (f_i^{(n)} u) = \begin{cases} q_i^{1-n} \frac{1 - q_i^{2(a+1-n)}}{1 - q_i^2} f_i^{(n-1)} u & \text{if } i \in I^{\text{re}}, \\ \frac{(1 - q_i^{2nc_i})(1 - q_i^{2(a+(n-1)c_i)})}{(1 - q_i^2)(1 - q_i^{2c_i})} f_i^{(n-1)} u & \text{if } a_{ii} < 0, \\ n \frac{1 - q_i^{2a}}{1 - q_i^2} f_i^{(n-1)} u & \text{if } a_{ii} = 0, \end{cases}$$

where $a = \lambda(h_i)$. Hence when $a_{ii} \leq 0$, we can use $q_i^{-1} K_i e_i$ instead of $\tilde{e}_i$.

Definition 4.6. Let M be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{\text{int}}$. A *crystal basis* of M is a pair (L, B) such that

- (i) L is a crystal lattice of M ,
- (ii) B is a $\mathbf{Q}$ -basis of $L/qL \cong \mathbf{Q} \otimes_{\mathbf{A}_0} L$,
- (iii) $B = \bigsqcup_{\lambda \in P} B_\lambda$, where $B_\lambda := B \cap (L_\lambda/qL_\lambda)$,
- (iv) $\tilde{e}_i B \subset B \cup \{0\}$ and $\tilde{f}_i B \subset B \cup \{0\}$ for every $i \in I$,
- (v) for any $i \in I$ and $b, b' \in B$, we have $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$.

The set B is endowed with a colored oriented graph structure whose arrows are defined by

$$b \xrightarrow{i} b' \quad \text{if and only if} \quad \tilde{f}_i b = b'.$$

The graph B is called the *crystal graph* of M and it reflects the combinatorial structure of M . For instance, we have

$$\dim_{\mathbf{Q}(q)} M_\lambda = \text{rank}_{\mathbf{A}_0} L_\lambda = \#B_\lambda \quad \text{for every } \lambda \in P.$$

(See, for example, [6, Theorem 4.2.5]. See also Proposition 4.9 below.)

Theorem 4.7. Let M be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{\text{int}}$ and let (L, B) be a crystal basis of M . For $i \in I$ and $u \in L_\lambda$, let $u = \sum_{l \geq 0} f_i^{(l)} u_l$ be the i -string decomposition of u . Then the following statements are true.

- (a) $u_l \in L$ for every $l \in \mathbf{Z}_{\geq 0}$.
- (b) If $u + qL \in B$, then there exists a non-negative integer l_0 such that
 - (i) $u_l \in qL$ for every $l \neq l_0$,
 - (ii) $u_{l_0} + qL \in B$,

$$(iii) \ u \equiv f_i^{(l_0)} u_{l_0} \pmod{qL}.$$

In particular, we have $\tilde{e}_i u \equiv f_i^{(l_0-1)} u_{l_0}$ and $\tilde{f}_i u \equiv f_i^{(l_0+1)} u_{l_0} \pmod{qL}$.

Proof. The proof is the same as the one given in [10, Proposition 2.3.2] (see also [6, Proposition 4.2.11]). $\square$

Let M be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ with a crystal basis (L, B) . For $i \in I$, we define the maps $\text{wt}: B \rightarrow P$, $\varepsilon_i: B \rightarrow \mathbf{Z}_{\geq 0}$ and $\varphi_i: B \rightarrow \mathbf{Z}_{\geq 0} \cup \{\infty\}$ by

$$(4.7) \quad \begin{aligned} \text{wt}(b) &= \lambda \quad \text{for } b \in B_\lambda \ (\lambda \in P), \\ \varepsilon_i(b) &= \max \{l \geq 0; \tilde{e}_i^l b \in B\}, \\ \varphi_i(b) &= \max \{l \geq 0; \tilde{f}_i^l b \in B\}. \end{aligned}$$

Note that for $i \in I^{\text{im}}$, we have (see Example 5.1)

$$\varphi_i(b) = \begin{cases} 0 & \text{if } \langle h_i, \text{wt}(b) \rangle = 0, \\ \infty & \text{if } \langle h_i, \text{wt}(b) \rangle > 0. \end{cases}$$

Then, using Theorem 4.7, we can easily prove the following proposition.

Proposition 4.8.

(a) If $b \in B_\lambda$ satisfies $\tilde{e}_i b \in B$, then

$$\text{wt}(\tilde{e}_i b) = \lambda + \alpha_i, \quad \varepsilon_i(\tilde{e}_i b) = \varepsilon_i(b) - 1, \quad \varphi_i(\tilde{e}_i b) = \varphi_i(b) + 1.$$

(b) If $b \in B_\lambda$ satisfies $\tilde{f}_i b \in B$, then

$$\text{wt}(\tilde{f}_i b) = \lambda - \alpha_i, \quad \varepsilon_i(\tilde{f}_i b) = \varepsilon_i(b) + 1, \quad \varphi_i(\tilde{f}_i b) = \varphi_i(b) - 1.$$

Proposition 4.9. For any $n \in \mathbf{Z}_{\geq 0}$ and $i \in I$, we have

$$(f_i^n M \cap L) / (f_i^n M \cap qL) = \bigoplus_{\substack{b \in B \\ \varepsilon_i(b) \geq n}} \mathbf{Q}b.$$

In particular, we have

$$\dim_{\mathbf{Q}(q)} (f_i^n M)_\lambda = \# \{b \in B_\lambda; \varepsilon_i(b) \geq n\}.$$

Proof. Since $L = \bigoplus_{m \geq 0} f_i^{(m)}(\text{Ker } e_i \cap L)$ by Lemma 4.5, one has

$$L/qL = \bigoplus_{m \geq 0} f_i^{(m)}(\text{Ker } e_i \cap L)/qf_i^{(m)}(\text{Ker } e_i \cap L).$$

If $b \in B$ satisfies $\varepsilon_i(b) = m$, then b belongs to $f_i^{(m)}(\text{Ker } e_i \cap L)/qf_i^{(m)}(\text{Ker } e_i \cap L)$. Hence $\{b \in B; \varepsilon_i(b) = m\}$ is a basis of $f_i^{(m)}(\text{Ker } e_i \cap L)/qf_i^{(m)}(\text{Ker } e_i \cap L)$. To complete the proof, it is enough to note that $f_i^n M \cap L = \bigoplus_{m \geq n} f_i^{(m)}(\text{Ker } e_i \cap L)$. $\square$

5. Tensor Product Rule

Fix $i \in I$, and let U_i be the subalgebra of $U_q(\mathfrak{g})$ generated by e_i, f_i and $U_q^0(\mathfrak{g})$. In the following example, we investigate the structure of crystal bases for irreducible highest weight U_i -modules.

Example 5.1. Let $V := V(\lambda)$ be the irreducible highest weight U_i -module with $m := \lambda(h_i) \in \mathbf{Z}_{\geq 0}$ and highest weight vector v_λ .

(a) If $a_{ii} = 2$, then we have

$$K_i v_\lambda = q_i^m v_\lambda, \quad e_i v_\lambda = 0, \quad f_i^{m+1} v_\lambda = 0,$$

and $V = \bigoplus_{l=0}^m \mathbf{Q}(q) f_i^{(l)} v_\lambda$. Set

$$L = \bigoplus_{l=0}^m \mathbf{A}_0 f_i^{(l)} v_\lambda \quad \text{and} \quad B = \{v_\lambda, f_i v_\lambda, \dots, f_i^{(m)} v_\lambda\}.$$

Then (L, B) is a crystal basis of V with the Kashiwara operators given by

$$\tilde{e}_i(f_i^{(l)} v_\lambda) = f_i^{(l-1)} v_\lambda, \quad \tilde{f}_i(f_i^{(l)} v_\lambda) = f_i^{(l+1)} v_\lambda.$$

Moreover, we have

$$\text{wt}(f_i^{(l)} v_\lambda) = \lambda - l\alpha_i, \quad \varepsilon_i(f_i^{(l)} v_\lambda) = l, \quad \varphi_i(f_i^{(l)} v_\lambda) = m - l.$$

(b) If $a_{ii} \leq 0$ and $\lambda(h_i) = 0$, then we have

$$K_i v_\lambda = v_\lambda, \quad e_i v_\lambda = f_i v_\lambda = 0, \quad \text{and} \quad V = \mathbf{Q}(q) v_\lambda.$$

Set $L = \mathbf{A}_0 v_\lambda$ and $B = \{v_\lambda\}$. Then (L, B) is a crystal basis of V , and

$$\tilde{e}_i v_\lambda = \tilde{f}_i v_\lambda = 0, \quad \text{wt}(v_\lambda) = \lambda, \quad \varepsilon_i(v_\lambda) = \varphi_i(v_\lambda) = 0.$$

(c) If $a_{ii} \leq 0$ and $m := \lambda(h_i) > 0$, then we have

$$K_i v_\lambda = q_i^m v_\lambda, \quad e_i v_\lambda = 0, \quad \text{and} \quad V = \bigoplus_{l \geq 0} \mathbf{Q}(q) f_i^l v_\lambda.$$

Set

$$L = \bigoplus_{l \geq 0} \mathbf{A}_0 f_i^l v_\lambda \quad \text{and} \quad B = \{f_i^l v_\lambda; l \geq 0\}.$$

Then (L, B) is a crystal basis of V with the Kashiwara operators given by

$$\tilde{e}_i(f_i^l v_\lambda) = f_i^{l-1} v_\lambda, \quad \tilde{f}_i(f_i^l v_\lambda) = f_i^{l+1} v_\lambda.$$

Moreover, we have

$$\text{wt}(f_i^l v_\lambda) = \lambda - l\alpha_i, \quad \varepsilon_i(f_i^l v_\lambda) = l, \quad \varphi_i(f_i^l v_\lambda) = \infty.$$

Now we are ready to state the *tensor product rule* for crystal bases of $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_{int}$.

Theorem 5.2. *Let M_j be a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_{int}$ and let (L_j, B_j) be a crystal basis of M_j ($j = 1, 2$). Set*

$$M = M_1 \otimes_{\mathbf{Q}(q)} M_2, \quad L = L_1 \otimes_{\mathbf{A}_0} L_2, \quad B = B_1 \otimes B_2.$$

Then (L, B) is a crystal basis of M , where the Kashiwara operators $\tilde{e}_i$ and $\tilde{f}_i$ ($i \in I$) on B are given as follows:

$$\begin{aligned} \tilde{e}_i(b_1 \otimes b_2) &= \begin{cases} \tilde{e}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) \geq \varepsilon_i(b_2), \\ b_1 \otimes \tilde{e}_i b_2 & \text{if } \varphi_i(b_1) < \varepsilon_i(b_2), \end{cases} \\ \tilde{f}_i(b_1 \otimes b_2) &= \begin{cases} \tilde{f}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) > \varepsilon_i(b_2), \\ b_1 \otimes \tilde{f}_i b_2 & \text{if } \varphi_i(b_1) \leq \varepsilon_i(b_2). \end{cases} \end{aligned}$$

Proof. By Theorem 4.7, it suffices to prove the tensor product rule for irreducible highest weight U_i -modules $M_1 = V(\lambda)$ and $M_2 = V(\mu)$ with $a := \lambda(h_i) \in \mathbf{Z}_{\geq 0}$ and $b := \mu(h_i) \in \mathbf{Z}_{\geq 0}$.

If $a_{ii} = 2$, our assertion was already proved in [9]. (See also [6, Theorem 4.4.3], [11, Théorème 2.3.5]).

If $a_{ii} \leq 0$, set $c_i = -\frac{1}{2}a_{ii} \in \mathbf{Z}_{\geq 0}$. If $a = 0$ or $b = 0$, then our assertion is trivial. We assume that $a, b > 0$. Then, by (4.5), we have $\tilde{e}_i \circ \tilde{f}_i = \text{id}_{M_1 \otimes M_2}$ and

$$\begin{aligned} M_1 &= V(\lambda) = \bigoplus_{l \geq 0} \mathbf{Q}(q) f_i^l u, & M_2 &= V(\mu) = \bigoplus_{l \geq 0} \mathbf{Q}(q) f_i^l v, \\ L_1 &= \bigoplus_{l \geq 0} \mathbf{A}_0 f_i^l u, & L_2 &= \bigoplus_{l \geq 0} \mathbf{A}_0 f_i^l v, \\ B_1 &= \{f_i^l u \mod qL_1; l \in \mathbf{Z}_{\geq 0}\}, & B_2 &= \{f_i^l v \mod qL_2; l \in \mathbf{Z}_{\geq 0}\}, \end{aligned}$$

where

$$e_i u = e_i v = 0, \quad K_i u = q_i^{\lambda(h_i)} u = q_i^a u, \quad K_i v = q_i^{\mu(h_i)} v = q_i^b v.$$

Consider the $\mathbf{A}_0$ -lattice

$$L = L_1 \otimes_{\mathbf{A}_0} L_2 = \bigoplus_{l, m \geq 0} \mathbf{A}_0(f_i^l u \otimes f_i^m v).$$

We have

$$\begin{aligned} f_i(f_i^l u \otimes f_i^m v) &= (f_i \otimes 1 + K_i \otimes f_i)(f_i^l u \otimes f_i^m v) \\ &= f_i^{l+1} u \otimes f_i^m v + q_i^{a+2lc_i} f_i^l u \otimes f_i^{m+1} v \\ &\equiv f_i^{l+1} u \otimes f_i^m v \mod qL. \end{aligned}$$

Therefore, we have $\tilde{f}_i L \subset L$ and

$$\tilde{f}_i(f_i^l u \otimes f_i^m v) \equiv \tilde{f}_i(f_i^l u) \otimes f_i^m v \mod qL.$$

On the other hand, we have (see Lemma 4.5 and (4.6))

$$\begin{aligned} q_i^{-1} K_i e_i(f_i^l u \otimes f_i^m v) &= (q_i^{-1} K_i e_i \otimes 1 + K_i \otimes q_i^{-1} K_i e_i)(f_i^l u \otimes f_i^m v) \\ &= (q_i^{-1} K_i e_i f_i^l u) \otimes f_i^m v + q_i^{a+2c_i l} f_i^l u \otimes q_i^{-1} K_i e_i f_i^m v \\ &\equiv (q_i^{-1} K_i e_i f_i^l u) \otimes f_i^m v \mod qL. \end{aligned}$$

Therefore, we have $\tilde{e}_i L \subset L$, and $\tilde{e}_i(u \otimes f_i^m v) \equiv 0 \pmod{qL}$. For $l > 0$, we have

$$\tilde{e}_i(f_i^l \otimes f_i^m v) \equiv \tilde{e}_i \tilde{f}_i(f_i^{l-1} \otimes f_i^m v) = f_i^{l-1} \otimes f_i^m v \pmod{qL}.$$

Hence $(L_1 \otimes L_2, B_1 \otimes B_2)$ is a crystal basis with the desired action of $\tilde{e}_i$ and $\tilde{f}_i$. $\square$

6. Crystal basis for $U_q^-(\mathfrak{g})$

In this section, we will define the notion of crystal basis for $U_q^-(\mathfrak{g})$ following [10]. Although [10] treated the Kac-Moody case, similar arguments can be applied to the generalized Kac-Moody algebra case with slight modifications, and we omit the details.

Fix $i \in I$. For any $P \in U_q^-(\mathfrak{g})$, there exist unique $Q, R \in U_q^-(\mathfrak{g})$ such that

$$(6.1) \quad e_i P - P e_i = \frac{K_i Q - K_i^{-1} R}{q_i - q_i^{-1}}.$$

We define the endomorphisms $e'_i, e''_i: U_q^-(\mathfrak{g}) \rightarrow U_q^-(\mathfrak{g})$ by

$$(6.2) \quad e'_i(P) = R \quad \text{and} \quad e''_i(P) = Q.$$

Then we have the following commutation relations:

$$(6.3) \quad e'_i f_j = q_i^{-a_{ij}} f_j e'_i + \delta_{ij}, \quad e''_i f_j = q_i^{a_{ij}} f_j e''_i + \delta_{ij} \quad \text{for any } i, j \in I.$$

Recall that $c_i := -a_{ii}/2$ and set

$$e_i'^{(l)} = \begin{cases} (e'_i)^l & \text{if } a_{ii} = 2, \\ \frac{(e'_i)^l}{\{l\}_i!} & \text{if } a_{ii} \leq 0. \end{cases}$$

By the commutation relation (6.3), we have

$$(6.4) \quad e_i'^{(n)} f_j^{(m)} = \begin{cases} \sum_{k=0}^n q_i^{-2nm+(n+m)k-k(k-1)/2} \begin{bmatrix} n \\ k \end{bmatrix}_i f_i^{(m-k)} e_i'^{(n-k)} & \text{if } i = j \text{ and } a_{ii} = 2, \\ \sum_{k=0}^m q_i^{-c_i(-2nm+(n+m)k-k(k-1)/2)} \left\{ \begin{matrix} m \\ k \end{matrix} \right\}_i f_i^{(m-k)} e_i'^{(n-k)} & \text{if } i = j \text{ and } a_{ii} \leq 0, \\ q_i^{-nma_{ij}} f_j^{(m)} e_i'^{(n)} & \text{if } i \neq j. \end{cases}$$

By a similar argument in [10], one can show that there exists a unique non-degenerate symmetric bilinear form on $U_q^-(\mathfrak{g})$ satisfying

$$(6.5) \quad (1, 1) = 1, \quad (f_i P, Q) = (P, e'_i Q) \quad \text{for all } i \in I.$$

Let $\star$ be the anti-involution of $U_q(\mathfrak{g})$ defined by

$$e_i^\star = e_i, \quad f_i^\star = f_i \quad \text{and} \quad (q^h)^\star = q^{-h} \quad \text{for all } i \in I, h \in P^\vee.$$

Then we have

$$(6.6) \quad (P^\star, Q^\star) = (P, Q) \quad \text{for any } P, Q \in U_q^-(\mathfrak{g}).$$

We define the following operator on $U_q^-(\mathfrak{g})$:

$$(6.7) \quad P_i = \sum_{n \geq 0} (-1)^n q_i^{c_i n(n-1)/2} f_i^{(n)} e_i'^{(n)}.$$

Then P_i is the projection operator to $\text{Ker}(e'_i)$ with respect to the decomposition $U_q^-(\mathfrak{g}) = \text{Ker}(e'_i) \oplus f_i U_q^-(\mathfrak{g})$. More precisely, we have the following result.

Proposition 6.1. *For each $i \in I$, every $u \in U_q^-(\mathfrak{g})$ can be uniquely expressed as*

$$(6.8) \quad u = \sum_{l \geq 0} f_i^{(l)} u_l,$$

where $e'_i u_l = 0$ for every $l \geq 0$ and $u_l = 0$ for $l \gg 0$. Moreover, we have

$$u_l = q_i^{-c_i l(l-1)/2} P_i e_i'^{(l)} u.$$

We call (6.8) the *i-string decomposition* of u .

Definition 6.2. For $i \in I$ and $u \in U_q^-(\mathfrak{g})$, let $u = \sum_{l \geq 0} f_i^{(l)} u_l$ be the *i-string decomposition*. Then the *Kashiwara operators* $\tilde{e}_i$ and $\tilde{f}_i$ on $U_q^-(\mathfrak{g})$ are defined by

$$(6.9) \quad \tilde{e}_i u = \sum_{l \geq 1} f_i^{(l-1)} u_l, \quad \tilde{f}_i u = \sum_{l \geq 0} f_i^{(l+1)} u_l.$$

Note that the left multiplication operator $f_i: U_q^-(\mathfrak{g}) \rightarrow U_q^-(\mathfrak{g})$ is injective. Hence we have

$$(6.10) \quad \tilde{e}_i \circ \tilde{f}_i = \text{id}_{U_q^-(\mathfrak{g})}.$$

Definition 6.3. A *crystal basis* of $U_q^-(\mathfrak{g})$ is a pair (L, B) , where

- (i) L is a free $\mathbf{A}_0$ -submodule of $U_q^-(\mathfrak{g})$ such that $U_q^-(\mathfrak{g}) \simeq \mathbf{Q}(q) \otimes_{\mathbf{A}_0} L$,
- (ii) B is a $\mathbf{Q}$ -basis of $L/qL \simeq \mathbf{Q} \otimes_{\mathbf{A}_0} L$,
- (iii) $\tilde{e}_i L \subset L$ and $\tilde{f}_i L \subset L$,
- (iv) $\tilde{e}_i B \subset B \cup \{0\}$ and $\tilde{f}_i B \subset B$,
- (v) For $b, b' \in B$, we have $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$.

Proposition 6.4. *Let (L, B) be a crystal basis of $U_q^-(\mathfrak{g})$. For $i \in I$ and $u \in L_{-\alpha}$, consider the i -string decomposition*

$$u = \sum_{l \geq 0} f_i^{(l)} u_l \quad \text{with} \quad e'_i u_l = 0.$$

Then the following statements are true.

- (a) $u_l \in L$ for every $l \geq 0$.
- (b) *If $u + qL \in B$, then there exists a non-negative integer l_0 such that*
 - (i) $u_l \in qL$ for every $l \neq l_0$,
 - (ii) $u_{l_0} + qL \in B$,
 - (iii) $u \equiv f_i^{(l_0)} u_{l_0} \pmod{qL}$.

7. Existence of Crystal Bases

In this section, we will prove the existence of crystal bases for the algebra $U_q^-(\mathfrak{g})$ and for $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_{int}$. Assuming the existence theorem, the uniqueness of crystal bases can be proved by the same argument as in [10] or [6, Section 5.2].

Let $\lambda \in P^+$ be a dominant integral weight and let $V(\lambda)$ be the irreducible highest weight $U_q(\mathfrak{g})$ -module with highest weight λ and highest weight vector v_λ . Let $L(\lambda)$ be the $\mathbf{A}_0$ -submodule of $V(\lambda)$ generated by the vectors of the form

$$\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda \quad (r \geq 0, i_1, \dots, i_r \in I)$$

and set

$$\begin{aligned} B(\lambda) &:= \left\{ \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda + qL(\lambda); r \geq 0, i_1, \dots, i_r \in I \right\} \setminus \{0\} \\ &\subset L(\lambda)/qL(\lambda). \end{aligned}$$

Similarly, we define $L(\infty)$ to be the free $\mathbf{A}_0$ -submodule of $U_q^-(\mathfrak{g})$ generated by the vectors of the form

$$\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1 \quad (r \geq 0, i_1, \dots, i_r \in I)$$

and set

$$B(\infty) := \left\{ \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1 + qL(\infty); r \geq 0, i_1, \dots, i_r \in I \right\} \subset L(\infty)/qL(\infty).$$

Our goal is to prove:

Theorem 7.1.

- (a) *The pair $(L(\lambda), B(\lambda))$ is a crystal basis of $V(\lambda)$.*
- (b) *The pair $(L(\infty), B(\infty))$ is a crystal basis of $U_q^-(\mathfrak{g})$.*

It is easy to see that $L(\lambda)$ has the weight space decomposition $L(\lambda) = \bigoplus_{\mu \in P} L(\lambda)_\mu$, where $L(\lambda)_\mu$ is the $\mathbf{A}_0$ -submodule of $L(\lambda)$ generated by the vectors of the form $\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda$ such that $\lambda - (\alpha_{i_1} + \cdots + \alpha_{i_r}) = \mu$. Similarly, $B(\lambda)$ has the weight set decomposition $B(\lambda) = \bigsqcup_{\mu \in P} B(\lambda)_\mu$. By the definition, we have, for $\alpha \in Q^+ \setminus \{0\}$

$$L(\lambda)_{\lambda-\alpha} = \sum_{i \in I} \tilde{f}_i (L(\lambda)_{\lambda-\alpha+\alpha_i}) \quad \text{and} \quad B(\lambda)_{\lambda-\alpha} = \left(\bigcup_{i \in I} \tilde{f}_i (B(\lambda)_{\lambda-\alpha+\alpha_i}) \right) \setminus \{0\}.$$

Since $V(\lambda)_\mu = \sum_i \tilde{f}_i V(\lambda)_{\mu+\alpha_i}$, we can easily see that $V(\lambda) \cong \mathbf{Q}(q) \otimes_{\mathbf{A}_0} L(\lambda)$.

Hence in order to prove (a), it remains to prove the following statements:

- (1) $\tilde{e}_i L(\lambda) \subset L(\lambda)$ for every $i \in I$,
- (2) $\tilde{e}_i B(\lambda) \subset B(\lambda) \cup \{0\}$ for every $i \in I$,
- (3) for all $i \in I$ and $b, b' \in B(\lambda)$, $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$,
- (4) $B(\lambda)$ is a $\mathbf{Q}$ -basis of $L(\lambda)/qL(\lambda)$.

Similarly, to prove (b), we need to prove:

- (1) $\tilde{e}_i L(\infty) \subset L(\infty)$ for every $i \in I$,
- (2) $\tilde{e}_i B(\infty) \subset B(\infty) \cup \{0\}$ for every $i \in I$,
- (3) for all $i \in I$ and $b, b' \in B(\infty)$, $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$,
- (4) $B(\infty)$ is a $\mathbf{Q}$ -basis of $L(\infty)/qL(\infty)$.

We will prove these statements using an interlocking induction argument on weights, called the *grand-loop argument*, employed in [10].

For $\lambda \in P^+$, let $\pi_\lambda: U_q^-(\mathfrak{g}) \rightarrow V(\lambda)$ be the $U_q^-(\mathfrak{g})$ -module homomorphism given by $P \mapsto Pv_\lambda$. For dominant integral weights $\lambda, \mu \in P^+$, there exist unique $U_q(\mathfrak{g})$ -module homomorphisms

$$\begin{aligned}\Phi_{\lambda,\mu}: V(\lambda + \mu) &\rightarrow V(\lambda) \otimes V(\mu), \\ \Psi_{\lambda,\mu}: V(\lambda) \otimes V(\mu) &\rightarrow V(\lambda + \mu)\end{aligned}$$

satisfying

$$\Phi_{\lambda,\mu}(v_{\lambda+\mu}) = v_\lambda \otimes v_\mu, \quad \Psi_{\lambda,\mu}(v_\lambda \otimes v_\mu) = v_{\lambda+\mu}.$$

It is clear that $\Psi_{\lambda,\mu} \circ \Phi_{\lambda,\mu} = \text{id}_{V(\lambda+\mu)}$.

Let $(\ , \)$ denote the symmetric bilinear form on irreducible highest weight $U_q(\mathfrak{g})$ -modules defined in (2.3), and let us define a symmetric bilinear form on $V(\lambda) \otimes V(\mu)$ by

$$(7.1) \quad (u_1 \otimes u_2, v_1 \otimes v_2) = (u_1, v_1)(u_2, v_2)$$

for $u_1, v_1 \in V(\lambda)$ and $u_2, v_2 \in V(\mu)$. Then this form on $V(\lambda) \otimes V(\mu)$ also satisfies

$$(7.2) \quad (q^h u, v) = (u, q^h v), \quad (f_i u, v) = (u, q_i^{-1} K_i e_i v), \quad (e_i u, v) = (u, q_i f_i K_i^{-1} v)$$

for all $i \in I$ and $u, v \in V(\lambda) \otimes V(\mu)$. Moreover, we have

$$(7.3) \quad (\Psi_{\lambda,\mu}(u), v) = (u, \Phi_{\lambda,\mu}(v))$$

for all $u \in V(\lambda) \otimes V(\mu)$ and $v \in V(\lambda + \mu)$.

Let $r \in \mathbf{Z}_{\geq 0}$ be a non-negative integer, and set

$$Q_+(r) := \{\alpha \in Q_+; |\alpha| \leq r\}.$$

For $\lambda, \mu \in P^+$ and $\alpha \in Q_+(r)$, we will prove that the following interlocking inductive statements are true, which would complete the proof of Theorem 7.1.

A(r) : $\tilde{e}_i L(\lambda)_{\lambda-\alpha} \subset L(\lambda)$ for every $i \in I$.

B(r) : $\tilde{e}_i B(\lambda)_{\lambda-\alpha} \subset B(\lambda) \cup \{0\}$ for every $i \in I$.

C(r) : For all $i \in I$, $b \in B(\lambda)_{\lambda-\alpha+\alpha_i}$ and $b' \in B(\lambda)_{\lambda-\alpha}$, we have

$$\tilde{f}_i b = b' \text{ if and only if } b = \tilde{e}_i b'.$$

D(r) : $B(\lambda)_{\lambda-\alpha}$ is a $\mathbf{Q}$ -basis of $L(\lambda)_{\lambda-\alpha}/qL(\lambda)_{\lambda-\alpha}$.

- E**(r) : $\Phi_{\lambda,\mu}(L(\lambda + \mu)_{\lambda+\mu-\alpha}) \subset L(\lambda) \otimes L(\mu)$.
F(r) : $\Psi_{\lambda,\mu}((L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha}) \subset L(\lambda + \mu)$.
G(r) : $\Psi_{\lambda,\mu}((B(\lambda) \otimes B(\mu))_{\lambda+\mu-\alpha}) \subset B(\lambda + \mu) \cup \{0\}$.
H(r) : $\tilde{e}_i L(\infty)_{-\alpha} \subset L(\infty)$ for every $i \in I$.
I(r) : $\tilde{e}_i B(\infty)_{-\alpha} \subset B(\infty) \cup \{0\}$ for every $i \in I$.
J(r) : For all $i \in I$ and $b \in B(\infty)_{-\alpha}$, if $\tilde{e}_i b \neq 0$, then $b = \tilde{f}_i \tilde{e}_i b$.
K(r) : $B(\infty)_{-\alpha}$ is a **Q**-basis of $L(\infty)_{-\alpha}/qL(\infty)_{-\alpha}$.
L(r) : $\pi_\lambda(L(\infty)_{-\alpha}) = L(\lambda)_{\lambda-\alpha}$.
M(r) : For any $P \in L(\infty)_{-\alpha+\alpha_i}$, we have

$$\tilde{f}_i(Pv_\lambda) \equiv (\tilde{f}_i P)v_\lambda \pmod{qL(\lambda)}.$$

- N**(r) : If $\bar{\pi}_\lambda : (L(\infty)/qL(\infty))_{-\alpha} \longrightarrow (L(\lambda)/qL(\lambda))_{\lambda-\alpha}$ is the induced homomorphism, we have

$$\{b \in B(\infty)_{-\alpha}; \bar{\pi}_\lambda(b) \neq 0\} \xrightarrow{\sim} B(\lambda)_{\lambda-\alpha}.$$

- O**(r) : If $b \in B(\infty)_{-\alpha}$ satisfies $\bar{\pi}_\lambda(b) \neq 0$, then we have $\tilde{e}_i \bar{\pi}_\lambda(b) = \bar{\pi}_\lambda(\tilde{e}_i b)$.

Note that the above statements are true for $r = 0$ and $r = 1$. From now on, we assume that $r \geq 2$ and the statements **A**($r - 1$), ..., **O**($r - 1$) are true.

In the sequel, for $\lambda \in P^+$, “ $\lambda \gg 0$ ” means “ $\lambda(h_i) \gg 0$ for all $i \in I$ ”. For instance, “for any $P \in L(\infty)$, we have $Pv_\lambda \in L(\lambda)$ for $\lambda \gg 0$ ” is paraphrased as “for any $P \in L(\infty)$, there exists a positive integer m such that $Pv_\lambda \in L(\lambda)$ whenever $\lambda \in P^+$ satisfies $\lambda(h_i) \geq m$ for all $i \in I$ ”.

Lemma 7.2. *Let $\alpha \in Q_+(r - 1)$ and $b \in B(\lambda)_{\lambda-\alpha}$. If $\tilde{e}_i b = 0$ for every i , then we have $\alpha = 0$ and $b = v_\lambda$.*

Proof. If $\alpha \neq 0$, then there exist i and $b' \in B(\lambda)_{\lambda-\alpha+\alpha_i}$ such that $b = \tilde{f}_i b'$. Then **C**($r - 1$) implies $\tilde{e}_i b = b' \neq 0$. $\square$

Lemma 7.3. *For $\alpha \in Q_+(r - 1)$, let $u = \sum_{n=0}^N f_i^{(n)} u_n \in V(\lambda)_{\lambda-\alpha}$ be the i -string decomposition, where $u_n \in \text{Ker } e_i \cap V(\lambda)_{\lambda-\alpha+n\alpha_i}$.*

- (a) *If $u \in L(\lambda)$, then $u_n \in L(\lambda)$ for every $n \geq 0$.*

- (b) If $u + qL(\lambda) \in B(\lambda)$, then there exists a non-negative integer n_0 such that $u \equiv f_i^{(n_0)} u_{n_0} \pmod{qL(\lambda)}$ and $u_n \in qL(\lambda)$ for $n \neq n_0$.

Proof. (a) We will use the induction on N . If $N = 0$, there is nothing to prove. If $N > 0$, $\mathbf{A}(r-1)$ implies $\tilde{e}_i u = \sum_{n=1}^N f_i^{(n-1)} u_n \in L(\lambda)$. By the induction hypothesis, $u_n \in L(\lambda)$ for every $n = 1, \dots, N$. Hence $u_0 = u - \sum_{n=1}^N \tilde{f}_i^n u_n \in L(\lambda)$, which proves (a).

(b) If $N = 0$, our assertion is trivial. If $N > 0$, by $\mathbf{B}(r-1)$, we have $\tilde{e}_i u + qL(\lambda) \in B(\lambda) \cup \{0\}$.

If $\tilde{e}_i u \in qL(\lambda)$, then (a) implies that $u_n \in qL(\lambda)$ for every $n \geq 1$, and $u \equiv u_0 \pmod{qL(\lambda)}$.

If $\tilde{e}_i u \notin qL(\lambda)$, then $\tilde{e}_i u + qL(\lambda) \in B(\lambda)$. By the induction hypothesis, there exists $n_0 \geq 1$ such that $\tilde{e}_i u \equiv f_i^{(n_0-1)} u_{n_0} \pmod{qL(\lambda)}$. Hence by $\mathbf{C}(r-1)$, we obtain

$$u \equiv \tilde{f}_i \tilde{e}_i u \equiv \tilde{f}_i f_i^{(n_0-1)} u_{n_0} = f_i^{(n_0)} u_{n_0} \pmod{qL(\lambda)}.$$

Therefore $u_n \in qL(\lambda)$ for $n \neq n_0$ by (a). $\square$

Lemma 7.4. For $\alpha \in Q_+(r-1)$, let $u = \sum_{n=0}^N f_i^{(n)} u_n \in U_q^-(\mathfrak{g})_{-\alpha}$ be the i -string decomposition.

- (a) If $u \in L(\infty)$, then $u_n \in L(\infty)$ for every $n \geq 0$.
(b) If $u + qL(\infty) \in B(\infty)$, then there exists a non-negative integer n_0 such that $u \equiv f_i^{(n_0)} u_{n_0} \pmod{qL(\infty)}$ and $u_n \in qL(\infty)$ for $n \neq n_0$.

Proof. The proof is similar to that of the preceding lemma. $\square$

For $\alpha \in Q_+(r-1)$ and $b \in B(\lambda)_{\lambda-\alpha}$, we set $\varepsilon_i(b) = \max\{n \geq 0; \tilde{e}_i^n b \neq 0\}$. We define

$$\varphi_i(b) = \begin{cases} \varepsilon_i(b) + \langle h_i, \text{wt}(b) \rangle & \text{if } i \in I^{\text{re}}, \\ 0 & \text{if } i \in I^{\text{im}} \text{ and } \langle h_i, \text{wt}(b) \rangle = 0, \\ \infty & \text{if } i \in I^{\text{im}} \text{ and } \langle h_i, \text{wt}(b) \rangle > 0. \end{cases}$$

Note that $\varphi_i(b) = 0$ implies $\tilde{f}_i b = 0$.

Lemma 7.5. Suppose $\alpha, \beta \in Q_+(r-1)$ and $b \in B(\lambda)_{\lambda-\alpha}$, $b' \in B(\mu)_{\mu-\beta}$.

(a) For each $i \in I$, we have

$$\begin{aligned}\tilde{f}_i(L(\lambda)_{\lambda-\alpha} \otimes L(\mu)_{\mu-\beta}) &\subset L(\lambda) \otimes L(\mu), \\ \tilde{e}_i(L(\lambda)_{\lambda-\alpha} \otimes L(\mu)_{\mu-\beta}) &\subset L(\lambda) \otimes L(\mu).\end{aligned}$$

(b) For each $i \in I$, the tensor product rule holds in $(L(\lambda) \otimes L(\mu))/q(L(\lambda) \otimes L(\mu))$:

$$\begin{aligned}\tilde{f}_i(b \otimes b') &= \begin{cases} \tilde{f}_i b \otimes b' & \text{if } \varphi_i(b) > \varepsilon_i(b'), \\ b \otimes \tilde{f}_i b' & \text{if } \varphi_i(b) \leq \varepsilon_i(b'), \end{cases} \\ \tilde{e}_i(b \otimes b') &= \begin{cases} \tilde{e}_i b \otimes b' & \text{if } \varphi_i(b) \geq \varepsilon_i(b'), \\ b \otimes \tilde{e}_i b' & \text{if } \varphi_i(b) < \varepsilon_i(b'). \end{cases}\end{aligned}$$

(c) For each $i \in I$, we have

$$\begin{aligned}\tilde{f}_i(B(\lambda)_{\lambda-\alpha} \otimes B(\mu)_{\mu-\beta}) &\subset (B(\lambda) \otimes B(\mu)) \cup \{0\}, \\ \tilde{e}_i(B(\lambda)_{\lambda-\alpha} \otimes B(\mu)_{\mu-\beta}) &\subset (B(\lambda) \otimes B(\mu)) \cup \{0\}.\end{aligned}$$

(d) If $\tilde{e}_i(b \otimes b') \neq 0$, then $b \otimes b' \equiv \tilde{f}_i \tilde{e}_i(b \otimes b')$.

(e) If $\tilde{e}_i(b \otimes b') = 0$ for every $i \in I$, then $b = v_\lambda$.

(f) For each $i \in I$, we have

$$\tilde{f}_i(b \otimes v_\mu) \equiv \tilde{f}_i b \otimes v_\mu \quad \text{or} \quad \tilde{f}_i b = 0.$$

(g) For any sequence of indices $i_1, \dots, i_r \in I$, we have

$$\tilde{f}_{i_1} \cdots \tilde{f}_{i_r}(v_\lambda \otimes v_\mu) \equiv (\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda) \otimes v_\mu \pmod{qL(\lambda) \otimes L(\mu)}$$

$$\text{or } \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda \in qL(\lambda).$$

Proof. (a) By Lemma 7.3(a), it suffices to prove the following statement: for $u \in \text{Ker}(e_i) \cap L(\lambda)_{\lambda-\alpha+n\alpha_i}$ and $v \in \text{Ker}(e_i) \cap L(\mu)_{\mu-\beta+m\alpha_i}$, we have

$$\begin{aligned}\tilde{f}_i(f_i^{(n)} u \otimes f_i^{(m)} v) &\in L(\lambda) \otimes L(\mu), \\ \tilde{e}_i(f_i^{(n)} u \otimes f_i^{(m)} v) &\in L(\lambda) \otimes L(\mu).\end{aligned}$$

Let L be the free $\mathbf{A}_0$ -submodule of $V(\lambda) \otimes V(\mu)$ generated by the vectors $f_i^{(s)} u \otimes f_i^{(t)} v$ ($s, t \geq 0$). Then by the tensor product rule for U_i -modules, we have $\tilde{e}_i L \subset L$

and $\tilde{f}_i L \subset L$. Hence our assertion follows immediately from the fact that $L \subset L(\lambda) \otimes L(\mu)$.

(b), (d) By Lemma 7.3(b), we may assume that $b = f_i^{(n)} u + qL(\lambda)$ and $b' = f_i^{(m)} v + qL(\mu)$ with $e_i u = 0$ and $e_i v = 0$. Let L be the $\mathbf{A}_0$ -submodule generated by $f_i^{(s)} u \otimes f_i^{(t)} v$ ($s, t \geq 0$). Then by the tensor product rule for U_i -modules, our assertion holds in L/qL . Since $L \subset L(\lambda) \otimes L(\mu)$, we are done.

The statements (c) is a simple consequence of (b).

(e) By (b), we have $\tilde{e}_i b = 0$ and Lemma 7.2 implies the desired result.

(f) By the tensor product rule (b), we have $\tilde{f}_i(b \otimes v_\mu) \equiv \tilde{f}_i b \otimes v_\mu$ unless $\varphi_i(b) = 0$, in which case $\tilde{f}_i b = 0$.

(g) This is an immediate consequence of (f). $\square$

Proposition 7.6 ($\mathbf{E}(r)$). *For every $\alpha \in Q_+(r)$, we have*

$$\Phi_{\lambda, \mu}(L(\lambda + \mu)_{\lambda + \mu - \alpha}) \subset L(\lambda) \otimes L(\mu).$$

Proof. Recall that $L(\lambda + \mu)_{\lambda + \mu - \alpha} = \sum_i \tilde{f}_i L(\lambda + \mu)_{\lambda + \mu - \alpha + \alpha_i}$. By $\mathbf{E}(r-1)$ and Lemma 7.5(a), we obtain the desired result. $\square$

Lemma 7.7. *Let $i_1, \dots, i_r$ be a sequence of indices in I . Suppose that $i_t \neq i_{t+1} = \dots = i_r$ for a positive integer $t < r$. Then for any dominant integral weight $\mu \in P^+$ and $\lambda = \Lambda_{i_t}$, we have*

$$\tilde{f}_{i_1} \cdots \tilde{f}_{i_r}(v_\lambda \otimes v_\mu) \equiv b \otimes b' \pmod{qL(\lambda) \otimes L(\mu)}$$

for some $b \in B(\lambda)_{\lambda - \alpha} \cup \{0\}$, $b' \in B(\mu)_{\mu - \beta} \cup \{0\}$ and $\alpha, \beta \in Q_+(r-1) \setminus \{0\}$.

Proof. Since $\Lambda_{i_t}(h_{i_r}) = 0$, we have $f_{i_r} v_\lambda = 0$. Hence we get

$$f_{i_r}(v_\lambda \otimes v) = (f_{i_r} \otimes 1 + K_{i_r} \otimes f_{i_r})(v_\lambda \otimes v) = v_\lambda \otimes f_{i_r} v$$

for every $v \in V(\mu)$, which gives

$$\tilde{f}_{i_{t+1}} \cdots \tilde{f}_{i_r}(v_\lambda \otimes v_\mu) = f_{i_r}^{(r-t)}(v_\lambda \otimes v_\mu) = v_\lambda \otimes f_{i_r}^{(r-t)} v_\mu.$$

Since $e_{i_t}(v_\lambda \otimes f_{i_r}^{(r-t)}v_\mu) = 0$, we have

$$\begin{aligned}
\tilde{f}_{i_t} \cdots \tilde{f}_{i_r}(v_\lambda \otimes v_\mu) &= \tilde{f}_{i_t}(v_\lambda \otimes f_{i_r}^{(r-t)}v_\mu) = f_{i_t}(v_\lambda \otimes f_{i_r}^{(r-t)}v_\mu) \\
&= (f_{i_t} \otimes 1 + K_{i_t} \otimes f_{i_t})(v_\lambda \otimes f_{i_r}^{(r-t)}v_\mu) \\
&= f_{i_t}v_\lambda \otimes f_{i_r}^{(r-t)}v_\mu + q_{i_t}^{\lambda(h_{i_t})}v_\lambda \otimes f_{i_t}f_{i_r}^{(r-t)}v_\mu \\
&= \tilde{f}_{i_t}v_\lambda \otimes \tilde{f}_{i_r}^{r-t}v_\mu + q_{i_t}v_\lambda \otimes \tilde{f}_{i_t}\tilde{f}_{i_r}^{r-t}v_\mu \\
&\equiv \tilde{f}_{i_t}v_\lambda \otimes \tilde{f}_{i_r}^{r-t}v_\mu \pmod{qL(\lambda) \otimes L(\mu)}.
\end{aligned}$$

Hence Lemma 7.5(b) yields our assertion. $\square$

Lemma 7.8. *For every $\alpha \in Q_+(r)$, we have*

$$(L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha} = \sum_i \tilde{f}_i(L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha+\alpha_i} + v_\lambda \otimes L(\mu)_{\mu-\alpha}.$$

Proof. Let L be the left-hand side and L' be the right-hand side of the above equation. We already know $L' \subset L$ by Lemma 7.5(a). By Lemma 7.5(e), for any $\beta \in Q_+(r-1) \setminus \{0\}$ and $b \otimes b' \in B(\lambda)_{\lambda-\beta} \otimes B(\mu)_{\mu-\alpha+\beta}$, there exists $i \in I$ such that $\tilde{e}_i(b \otimes b') \neq 0$. By Lemma 7.5(d), we have

$$b \otimes b' \equiv \tilde{f}_i \tilde{e}_i(b \otimes b') \pmod{qL(\lambda) \otimes L(\mu)}.$$

Hence $L(\lambda)_{\lambda-\beta} \otimes L(\mu)_{\mu-\alpha+\beta} \subset L' + qL$, and we obtain

$$\begin{aligned}
L &= L(\lambda)_{\lambda-\alpha} \otimes v_\mu + \sum_{\beta \in Q_+(r-1) \setminus \{0\}} L(\lambda)_{\lambda-\beta} \otimes L(\mu)_{\mu-\alpha+\beta} + v_\lambda \otimes L(\mu)_{\mu-\alpha} \\
&\subset L(\lambda)_{\lambda-\alpha} \otimes v_\mu + L' + qL.
\end{aligned}$$

Note that for any $\tilde{f}_{i_1} \cdots \tilde{f}_{i_r}v_\lambda \in B(\lambda)_{\lambda-\alpha}$, Lemma 7.5(f) yields

$$(\tilde{f}_{i_1} \cdots \tilde{f}_{i_r}v_\lambda) \otimes v_\mu \equiv \tilde{f}_{i_1}\tilde{f}_{i_2} \cdots \tilde{f}_{i_r}(v_\lambda \otimes v_\mu) \pmod{qL(\lambda) \otimes L(\mu)}.$$

It follows that $L(\lambda)_{\lambda-\alpha} \otimes v_\mu \subset L' + qL$ and hence $L \subset L' + qL$. Then by Nakayama's Lemma, we conclude that $L = L'$. $\square$

For $\lambda, \mu \in P^+$, define a linear map $S_{\lambda, \mu}: V(\lambda) \otimes V(\mu) \rightarrow V(\lambda)$ by

$$(7.4) \quad \begin{aligned} S_{\lambda, \mu}(u \otimes v_\mu) &= u \quad \text{for every } u \in V(\lambda), \\ S_{\lambda, \mu}(V(\lambda) \otimes \sum_i f_i V(\mu)) &= 0. \end{aligned}$$

Note that $S_{\lambda, \mu}$ is a $U_q^-(\mathfrak{g})$ -module homomorphism.

Lemma 7.9.

- (a) For all $\lambda, \mu \in P^+$, we have $S_{\lambda, \mu}(L(\lambda) \otimes L(\mu)) = L(\lambda)$.
- (b) For all $\alpha \in Q_+(r-1)$ and $w \in (L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha}$, we have

$$S_{\lambda, \mu} \circ \tilde{f}_i(w) \equiv \tilde{f}_i \circ S_{\lambda, \mu}(w) \pmod{qL(\lambda)}.$$

Proof. (a) is obvious.

(b) By Lemma 7.3, we may assume that $w = f_i^{(n)}u \otimes f_i^{(m)}v$ with $u \in L(\lambda)$, $v \in L(\mu)$ and $e_i u = e_i v = 0$. Let L be the $\mathbf{A}_0$ -submodule generated by $f_i^{(s)}u \otimes f_i^{(t)}v$ ($s, t \geq 0$). Then the tensor product rule for U_i -modules yields

$$\tilde{f}_i w \equiv f_i^{(n+1)}u \otimes f_i^{(m)}v \quad \text{or} \quad f_i^{(n)}u \otimes f_i^{(m+1)}v \pmod{qL}.$$

Observe that $S_{\lambda, \mu}(\tilde{f}_i w) \equiv \tilde{f}_i(S_{\lambda, \mu}w) \equiv 0 \pmod{qL}$ unless $m = 0$ and $v \in L(\mu)_\mu = \mathbf{A}_0 v_\mu$. If $m = 0$ and $v = v_\mu$, then we have

$$\tilde{f}_i w = \tilde{f}_i(f_i^{(n)}u \otimes v_\mu) \equiv \begin{cases} f_i^{(n+1)}u \otimes v_\mu \pmod{qL} & \text{if } f_i^{(n+1)}u \neq 0, \\ f_i^{(n)}u \otimes f_i v_\mu \pmod{qL} & \text{if } f_i^{(n+1)}u = 0. \end{cases}$$

In both cases, we have $S_{\lambda, \mu}(\tilde{f}_i w) \equiv f_i^{(n+1)}u = \tilde{f}_i f_i^{(n)}u = \tilde{f}_i S_{\lambda, \mu}(w) \pmod{qL(\lambda)}$. $\square$

Lemma 7.10. Let $\alpha \in Q_+(r)$ and $m \in \mathbf{Z}_{\geq 0}$. Then there exists a positive integer N satisfying the following properties: if $\lambda \in P^+$ and $u \in V(\lambda)_{\lambda-\alpha}$ satisfies $q_i^{-1}K_i e_i u \in q^N L(\lambda)$, then we have

- (a) $\tilde{f}_i^m u \equiv f_i^{(m)}u \pmod{qL(\lambda)}$,
- (b) $\tilde{e}_i f_i^{(m)}u \equiv f_i^{(m-1)}u \pmod{qL(\lambda)}$ if $i \in I^{\text{re}}$ and $m \leq \langle h_i, \lambda - \alpha \rangle$ or if $i \in I^{\text{im}}$ and $\lambda(h_i) > 0$.

Proof. Let $u = \sum_{n=0}^r f_i^{(n)} u_n$ be the i -string decomposition of u . Then we have

$$q_i^{-1} K_i e_i u = \sum_{n=1}^r q_i^{-1} K_i e_i f_i^{(n)} u_n$$

with $q_i^{-1} K_i e_i f_i^{(n)} u_n \in f_i^{(n-1)}(\text{Ker } e_i)$. Hence Lemma 7.3 implies that $q_i^{-1} K_i e_i f_i^{(n)} u_n \in q^N L(\lambda)$ for $n \geq 1$. By (4.6), we have $f_i^{(n-1)} u_n \in \mathbf{A}_0(q_i^{-1} K_i e_i f_i^{(n)} u_n)$ for $n \geq 1$. We then conclude $u_n \in q^N L(\lambda)$ for $n \geq 1$ by Lemma 7.3.

Hence if $N \gg 0$, the elements $f_i^{(m)} f_i^{(n)} u_n$, $f_i^{(m-1)} f_i^{(n)} u_n$, $f_i^{(m+n)} u_n$ belong to $qL(\lambda)$ for $n \geq 1$, which implies

$$\tilde{f}_i^m u = \sum_{n=0}^r f_i^{(n+m)} u_n \equiv f_i^{(m)} u_0 \pmod{qL(\lambda)}$$

and

$$f_i^{(m)} u = \sum_{n=0}^r f_i^{(m)} f_i^{(n)} u_n \equiv f_i^{(m)} u_0 \pmod{qL(\lambda)}.$$

Hence we obtain $\tilde{f}_i^m u \equiv f_i^{(m)} u$. Similarly, we have $\tilde{f}_i^{m-1} u = f_i^{(m-1)} u \equiv f_i^{(m-1)} u_0$.

On the other hand, we have $f_i^{(m)} u = \sum_{n=0}^r a_n f_i^{(m+n)} u_n$, where a_n is a q -binomial coefficient or 1 according as i is real or imaginary. Since $a_n f_i^{(n+m-1)} u_n \in a_n q^N L(\lambda) \subset qL(\lambda)$ for $n \geq 1$ and $N \gg 0$, (4.4) yields

$$\begin{aligned} \tilde{e}_i f_i^{(m)} u &= \sum_{f_i^{(m+n)} u_n \neq 0} a_n f_i^{(n+m-1)} u_n \\ &\equiv \begin{cases} f_i^{(m-1)} u_0 & \text{if } f_i^{(m)} u_0 \neq 0 \pmod{qL(\lambda)}, \\ 0 & \text{if } f_i^{(m)} u_0 = 0 \pmod{qL(\lambda)}. \end{cases} \end{aligned}$$

However, by the conditions in (b), $f_i^{(m)} u_0 = 0$ implies $u_0 = 0$. Therefore we conclude $\tilde{e}_i f_i^{(m)} u \equiv f_i^{(m-1)} u_0 \equiv f_i^{(m-1)} u \pmod{qL(\lambda)}$. $\square$

Lemma 7.11. *Let $\alpha \in Q_+(r)$ and $P \in U_q^-(\mathfrak{g})_{-\alpha}$. Then for every $\lambda \gg 0$, we have*

$$\begin{cases} (\tilde{f}_i P) v_\lambda \equiv \tilde{f}_i (P v_\lambda) \pmod{qL(\lambda)}, \\ (\tilde{e}_i P) v_\lambda \equiv \tilde{e}_i (P v_\lambda) \pmod{qL(\lambda)}. \end{cases}$$

Proof. We may assume that $P = f_i^{(n)}Q$ with $e'_i Q = 0$. Then

$$\begin{aligned} q_i^{-1} K_i e_i (Q u_\lambda) &= (1 - q_i^2)^{-1} (e'_i Q - K_i^2 e''_i Q) u_\lambda \\ &= -(1 - q_i^2)^{-1} q_i^{2\langle h_i, \lambda + \alpha_i + \text{wt}(Q) \rangle} (e''_i Q) u_\lambda \end{aligned}$$

Hence by the preceding lemma, we have for $\lambda \gg 0$

$$\tilde{f}_i(f_i^{(n)} Q v_\lambda) \equiv f_i^{(n+1)} Q v_\lambda \quad \text{and} \quad \tilde{e}_i(f_i^{(n)} Q v_\lambda) \equiv f_i^{(n-1)} Q v_\lambda.$$

□

Proposition 7.12 ($\mathbf{M}(r)$). *For any $\lambda \in P^+$, $\alpha \in Q_+(r-1)$ and $P \in L(\infty)_{-\alpha}$, we have*

$$(\tilde{f}_i P) v_\lambda \equiv \tilde{f}_i(P v_\lambda) \pmod{qL(\lambda)}.$$

In particular, we have $(\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1) v_\lambda \equiv \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda \pmod{qL(\lambda)}$.

Proof. Take $\mu \gg 0$. Since $\lambda + \mu \gg 0$, Lemma 7.11 yields

$$(\tilde{f}_i P) v_{\lambda+\mu} \equiv \tilde{f}_i(P v_{\lambda+\mu}) \pmod{qL(\lambda + \mu)}.$$

By Proposition 7.6, applying $\Phi_{\lambda, \mu}$ gives

$$(7.5) \quad (\tilde{f}_i P)(v_\lambda \otimes v_\mu) \equiv \tilde{f}_i(P(v_\lambda \otimes v_\mu)) \pmod{qL(\lambda) \otimes L(\mu)}.$$

On the other hand, we have

$$P(v_\lambda \otimes v_\mu) = \Phi_{\lambda, \mu}(P v_{\lambda+\mu}) \in L(\lambda) \otimes L(\mu)$$

by $\mathbf{L}(r-1)$ and $\mathbf{E}(r-1)$. Applying $S_{\lambda, \mu}$ to (7.5), Lemma 7.9 implies

$$(\tilde{f}_i P) v_\lambda \equiv \tilde{f}_i(P v_\lambda) \pmod{qL(\lambda)},$$

as desired. □

Proposition 7.13 ($\mathbf{L}(r)$). *For any $\lambda \in P^+$ and $\alpha \in Q_+(r)$, we have*

$$\pi_\lambda(L(\infty)_{-\alpha}) = L(\lambda)_{\lambda-\alpha}.$$

Proof. By Proposition 7.12, we have

$$\pi_\lambda(\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1) \equiv \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda \pmod{qL(\lambda)}.$$

It follows that

$$\pi_\lambda(L(\infty)_{-\alpha}) \subset L(\lambda)_{\lambda-\alpha} \text{ and } L(\lambda)_{\lambda-\alpha} \subset \pi_\lambda(L(\infty)_{-\alpha}) + qL(\lambda)_{\lambda-\alpha}.$$

Now the desired result follows from Nakayama's Lemma. $\square$

Corollary 7.14. *Let $\bar{\pi}_\lambda: (L(\infty)/qL(\infty))_{-\alpha} \longrightarrow (L(\lambda)/qL(\lambda))_{\lambda-\alpha}$ be the surjective $\mathbf{Q}$ -linear map induced by π_λ .*

(a) *For $\beta \in Q_+(r-1)$ and $b \in B(\infty)_{-\beta}$, we have*

$$\bar{\pi}_\lambda(\tilde{f}_i b) = \tilde{f}_i \bar{\pi}_\lambda(b).$$

(b) *For $\alpha \in Q_+(r)$ and $\lambda \in P^+$, we have*

$$(\bar{\pi}_\lambda B(\infty)_{-\alpha}) \setminus \{0\} = B(\lambda)_{\lambda-\alpha}.$$

(c) *Given $\alpha \in Q_+(r)$, take $\lambda \gg 0$. Then π_λ induces the isomorphisms*

$$L(\infty)_{-\alpha} \xrightarrow{\sim} L(\lambda)_{\lambda-\alpha}, \quad B(\infty)_{-\alpha} \setminus \{0\} \xrightarrow{\sim} B(\lambda)_{\lambda-\alpha}.$$

Proof. (a) is an immediate consequence of Proposition 7.12, and (b) follows from (a). Finally, (c) follows from the fact that $\pi_\lambda: U_q^-(\mathfrak{g})_{-\alpha} \rightarrow V(\lambda)_{\lambda-\alpha}$ is an isomorphism for $\lambda \gg 0$. $\square$

Now we will prove the statements **A**(r) and **H**(r). Fix $\lambda \in P^+$, $i \in I$ and $\alpha = \alpha_{i_1} + \cdots + \alpha_{i_r} \in Q_+(r)$. Let T be a finite subset of P^+ containing λ and the fundamental weights $\Lambda_{i_1}, \dots, \Lambda_{i_r}$. Since T is a finite set, we can take a sufficiently large $N_1 \geq 0$ such that

$$\tilde{e}_i L(\mu)_{\mu-\alpha} \subset q^{-N_1} L(\mu) \quad \text{for every } \mu \in T.$$

Let $N_2 \geq 0$ be a non-negative integer such that

$$\tilde{e}_i L(\infty)_{-\alpha} \subset q^{-N_2} L(\infty).$$

Then for every $\mu \gg 0$, by Lemma 7.11 and Proposition 7.13, we have

$$\begin{aligned}\tilde{e}_i L(\mu)_{\mu-\alpha} &= \tilde{e}_i(L(\infty)_{-\alpha} v_\mu) \subset (\tilde{e}_i L(\infty)_{-\alpha}) v_\mu + q L(\mu)_{\mu-\alpha} \\ &\subset q^{-N_2} L(\infty)_{-\alpha} v_\mu + q L(\mu)_{\mu-\alpha} \subset q^{-N_2} L(\mu).\end{aligned}$$

Therefore, given $\alpha \in Q_+(r)$, there exists a non-negative integer $N \geq 0$ such that

$$(7.6) \quad \begin{aligned}\tilde{e}_i L(\mu)_{\mu-\alpha} &\subset q^{-N} L(\mu) \quad \text{for every } \mu \gg 0, \\ \tilde{e}_i L(\tau)_{\tau-\alpha} &\subset q^{-N} L(\tau) \quad \text{for every } \tau \in T, \\ \tilde{e}_i L(\infty)_{-\alpha} &\subset q^{-N} L(\infty).\end{aligned}$$

Lemma 7.15. *Given $\alpha \in Q_+(r)$, let $N \geq 0$ be a non-negative integer satisfying (7.6). Then for every $\mu \gg 0$ and $\tau \in T$, we have*

$$\tilde{e}_i((L(\tau) \otimes L(\mu))_{\tau+\mu-\alpha}) \subset q^{-N}(L(\tau) \otimes L(\mu)).$$

Proof. For $u \in L(\tau)_{\tau-\beta}$ and $v \in L(\mu)_{\mu-\gamma}$ with $\alpha = \beta + \gamma$, let us show $\tilde{e}_i(u \otimes v) \in q^{-N}(L(\tau) \otimes L(\mu))$. If $\beta \neq 0$ and $\gamma \neq 0$, we already proved our assertion in Lemma 7.5(a).

If $\beta = 0$, then $\gamma = \alpha$ and we may assume $u = v_\tau$. Let $v = \sum_m f_i^{(m)} v_m$ with $e_i v_m = 0$ be the i -string decomposition of v . By (7.6), we have

$$\tilde{e}_i v = \sum_{m \geq 1} f_i^{(m-1)} v_m \in q^{-N} L(\mu).$$

Lemma 7.3 implies that $v_m \in q^{-N} L(\mu)$ for every $m \geq 1$. Let L be the $\mathbf{A}_0$ -submodule generated by the vectors $f_i^{(s)} v_\tau \otimes f_i^{(t)} v_m$ ($s, t \geq 0, m \geq 1$). Then we have $\tilde{e}_i L \subset L$. It follows that

$$\tilde{e}_i(v_\tau \otimes v) = \sum_{m \geq 1} \tilde{e}_i(v_\tau \otimes f_i^{(m)} v_m) \in L \subset q^{-N} L(\tau) \otimes L(\mu).$$

If $\beta = \alpha$ and $\gamma = 0$, a similar argument yields $\tilde{e}_i(u \otimes v_\mu) \in q^{-N} L(\tau) \otimes L(\mu)$. $\square$

Lemma 7.16. *Given $\alpha \in Q_+(r)$, let $N \geq 1$ be a positive integer satisfying (7.6). Then we have*

- (a) $\tilde{e}_i L(\mu)_{\mu-\alpha} \subset q^{-N+1} L(\mu)$ for all $\mu \gg 0$,
- (b) $\tilde{e}_i L(\tau)_{\tau-\alpha} \subset q^{-N+1} L(\tau)$ for every $\tau \in T$,

$$(c) \tilde{e}_i L(\infty)_{-\alpha} \subset q^{-N+1} L(\infty).$$

Proof. (a) Let $u = \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\mu \in L(\mu)_{\mu-\alpha}$. Suppose $i_1 = \cdots = i_r$. If $i = i_1$, then $\tilde{e}_i u = \tilde{e}_i f_i^{(r)} v_\mu = f_i^{(r-1)} v_\mu \in L(\mu)$, and if $i \neq i_1$, $\tilde{e}_i u = 0$. Hence we may assume that there exists a positive integer $t < r$ such that $i_t \neq i_{t+1} = \cdots = i_r$.

Suppose $\mu \gg 0$ and set $\lambda_0 = \Lambda_{i_t}$, $\mu' = \mu - \lambda_0 \gg 0$. By Lemma 7.7, we have

$$w := \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} (v_{\lambda_0} \otimes v_{\mu'}) \equiv v \otimes v' \pmod{qL(\lambda_0) \otimes L(\mu')}$$

for some $v \in L(\lambda_0)_{\lambda_0-\beta}$, $v' \in L(\mu')_{\mu'-\gamma}$, $\beta, \gamma \in Q_+(r-1) \setminus \{0\}$, $\alpha = \beta + \gamma$. Hence Lemma 7.5(a) and Lemma 7.15 yield

$$\begin{aligned} \tilde{e}_i w &\in L(\lambda_0) \otimes L(\mu') + q\tilde{e}_i(L(\lambda_0) \otimes L(\mu'))_{\lambda_0+\mu'-\alpha} \\ &\subset L(\lambda_0) \otimes L(\mu') + q^{-N+1}L(\lambda_0) \otimes L(\mu') = q^{-N+1}L(\lambda_0) \otimes L(\mu'). \end{aligned}$$

Namely, we have

$$\tilde{e}_i w \in q^{-N+1}(L(\lambda_0) \otimes L(\mu'))_{\lambda_0+\mu'-\alpha+\alpha_i}.$$

Therefore, applying $\Psi_{\lambda_0, \mu'}$, the statement $\mathbf{F}(r-1)$ yields

$$\tilde{e}_i u = \tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\mu \in q^{-N+1} L(\mu).$$

(b) Let $\tau \in T$ and $u = \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\tau \in L(\tau)_{\tau-\alpha}$. If $u \in qL(\tau)$, then our assertion follows immediately from (7.6). If $u \notin qL(\tau)$, then for any $\mu \in P^+$, Lemma 7.5(g) gives

$$(7.7) \quad \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} (v_\tau \otimes v_\mu) \equiv u \otimes v_\mu \pmod{qL(\tau) \otimes L(\mu)}.$$

If $\mu \gg 0$, then (a) implies

$$\tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_{\tau+\mu} \in q^{-N+1} L(\tau + \mu).$$

Applying $\Phi_{\tau, \mu}$, the statement $\mathbf{E}(r-1)$ gives

$$\tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} (v_\tau \otimes v_\mu) \in q^{-N+1} L(\tau) \otimes L(\mu).$$

Along with (7.7), Lemma 7.15 yields

$$\begin{aligned} (7.8) \quad \tilde{e}_i (u \otimes v_\mu) &\in q^{-N+1} L(\tau) \otimes L(\mu) + q\tilde{e}_i(L(\tau) \otimes L(\mu)) \\ &\subset q^{-N+1} L(\tau) \otimes L(\mu). \end{aligned}$$

Let $u = \sum_{m \geq 0} f_i^{(m)} u_m$ be the i -string decomposition of u . Then $\tilde{e}_i u = \sum_{m \geq 1} f_i^{(m-1)} u_m \in q^{-N} L(\tau)$ implies that $u_m \in q^{-N} L(\tau)$ for $m \geq 1$. Let L be the $\mathbf{A}_0$ -submodule generated by $f_i^{(s)} v_m \otimes f_i^{(t)} v_\mu$ ($s, t \geq 0, m \geq 1$). Then $L \subset q^{-N} L(\tau) \otimes L(\mu)$ and by the tensor product rule for U_i -modules, we obtain

$$\begin{aligned} \tilde{e}_i(u \otimes v_\mu) &= \sum_{m \geq 1} \tilde{e}_i(f_i^{(m)} u_m \otimes v_\mu) \\ &\equiv \sum_{m \geq 1} f_i^{(m-1)} u_m \otimes v_\mu = \tilde{e}_i u \otimes v_\mu \pmod{qL}. \end{aligned}$$

Hence we have $\tilde{e}_i(u \otimes v_\mu) \equiv \tilde{e}_i u \otimes v_\mu \pmod{q^{1-N} L(\tau) \otimes L(\mu)}$. Together with (7.8), we obtain $\tilde{e}_i u \otimes v_\mu \in q^{-N+1} L(\tau) \otimes L(\mu)$, which implies the desired result $\tilde{e}_i u \in q^{-N+1} L(\tau)$.

(c) Let $P \in L(\infty)_{-\alpha}$ and take $\mu \gg 0$. By Lemma 7.11, we have

$$(\tilde{e}_i P) v_\mu \equiv \tilde{e}_i(P v_\mu) \pmod{qL(\mu)}.$$

By Proposition 7.13, we get

$$\tilde{e}_i(P v_\mu) \in \tilde{e}_i L(\mu)_{\mu-\alpha} \subset q^{-N+1} L(\mu),$$

which implies $(\tilde{e}_i P) v_\mu \in q^{-N+1} L(\mu)$. Hence by Corollary 7.14 (c), we obtain $\tilde{e}_i P \in q^{-N+1} L(\infty)$. $\square$

Proposition 7.17 ($\mathbf{A}(r), \mathbf{H}(r)$). *For all $\lambda \in P^+$ and $\alpha \in Q_+(r)$, we have*

$$\tilde{e}_i L(\infty)_{-\alpha} \subset L(\infty), \quad \tilde{e}_i L(\lambda)_{\lambda-\alpha} \subset L(\lambda).$$

Proof. By applying Lemma 7.16 repeatedly, we conclude that $N = 0$, and our assertion follows immediately. $\square$

Corollary 7.18. *For $\alpha \in Q_+(r)$, $B(\infty)_{-\alpha}$ does not contain 0.*

Proof. For $b \in B(\infty)_{-\alpha}$, there exists $i \in I$ and $b' \in B(\infty)_{-\alpha+\alpha_i}$ such that $b = \tilde{f}_i b'$. Then $b' = \tilde{e}_i b$ does not vanish by $\mathbf{K}(r-1)$, and hence $b \neq 0$. $\square$

Lemma 7.19. *For $\alpha \in Q_+(r)$, $i \in I$, $\lambda \gg 0$ and $b \in B(\infty)_{-\alpha}$, we have*

$$\bar{\pi}_\lambda(\tilde{e}_i b) = \tilde{e}_i \bar{\pi}_\lambda(b).$$

Proof. This follows immediately from Lemma 7.11. $\square$

Since Lemma 7.5 depends only on $\mathbf{A}(r-1)$, we have:

Corollary 7.20. *Let $\lambda, \mu \in P^+$ and $\alpha, \beta \in Q_+(r)$.*

- (a) *If $u = \sum_{n \geq 0} f_i^{(n)} u_n \in L(\lambda)_{\lambda-\alpha}$ is the i -string decomposition of u , then $u_n \in L(\lambda)$ for every $n \geq 0$.*
- (b) *For each $i \in I$, we have*

$$\begin{aligned}\tilde{f}_i(L(\lambda)_{\lambda-\alpha} \otimes L(\mu)_{\mu-\beta}) &\subset L(\lambda) \otimes L(\mu), \\ \tilde{e}_i(L(\lambda)_{\lambda-\alpha} \otimes L(\mu)_{\mu-\beta}) &\subset L(\lambda) \otimes L(\mu).\end{aligned}$$

In order to prove the statements $\mathbf{B}(r)$ and $\mathbf{O}(r)$, we first prove:

Lemma 7.21. *Let $\lambda, \mu \in P^+$ and $\alpha \in Q_+(r)$. Then for every $u \in L(\lambda)_{\lambda-\alpha}$, we have*

$$\tilde{e}_i(u \otimes v_\mu) \equiv (\tilde{e}_i u) \otimes v_\mu \pmod{qL(\lambda) \otimes L(\mu)}.$$

Proof. By Corollary 7.20(a), we may assume $u = f_i^{(n)} w \neq 0$ with $e_i w = 0$ and $w \in L(\lambda)_{\lambda-\alpha+n\alpha_i}$. Let L be the $\mathbf{A}_0$ -submodule generated by $f_i^{(s)} w \otimes f_i^{(t)} v_\mu$ ($s, t \geq 0$). Then the tensor product rule for U_i -modules gives

$$\tilde{e}_i(f_i^{(n)} w \otimes v_\mu) \equiv f_i^{(n-1)} w \otimes v_\mu \pmod{qL}.$$

Our assertion immediately follows from $L \subset L(\lambda) \otimes L(\mu)$. $\square$

Proposition 7.22 ($\mathbf{I}(r)$). *For every $\alpha \in Q_+(r)$, we have*

$$\tilde{e}_i B(\infty)_{-\alpha} \subset B(\infty) \cup \{0\}.$$

Proof. Let $P = \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1 \in L(\infty)_{-\alpha}$ and $b = \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1 + qL(\infty) \in B(\infty)_{-\alpha}$. If $i_1 = \cdots = i_r$, our assertion is true as we have seen in the proof of Lemma 7.16 (a).

Hence we may assume that there exists a positive integer $t < r$ such that $i_t \neq i_{t+1} = \cdots = i_r$. Take $\mu \gg 0$ and set $\lambda_0 = \Lambda_{i_t}$, $\lambda = \lambda_0 + \mu$. Then Lemma 7.7 yields

$$\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} (v_{\lambda_0} \otimes v_\mu) \equiv v \otimes v' \pmod{qL(\lambda_0) \otimes L(\mu)}$$

for some $v \in L(\lambda_0)_{\lambda_0 - \beta}$, $v' \in L(\mu)_{\mu - \gamma}$ and $\beta, \gamma \in Q_+(r-1) \setminus \{0\}$ with $\alpha = \beta + \gamma$ such that $v + qL(\lambda_0) \in B(\lambda_0) \cup \{0\}$ and $v' + qL(\mu) \in B(\mu) \cup \{0\}$. Then the tensor product rule for U_i -modules gives

$$\begin{aligned} \tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} (v_{\lambda_0} \otimes v_\mu) &\equiv \tilde{e}_i (v \otimes v') \\ &\equiv \tilde{e}_i v \otimes v' \text{ or } v \otimes \tilde{e}_i v' \pmod{qL(\lambda_0) \otimes L(\mu)}. \end{aligned}$$

By $\mathbf{B}(r-1)$, we have

$$\tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} (v_{\lambda_0} \otimes v_\mu) + qL(\lambda_0) \otimes L(\mu) \in (B(\lambda_0) \otimes B(\mu)) \cup \{0\}.$$

Hence, by applying $\Psi_{\lambda_0, \mu}$, the statement $\mathbf{G}(r-1)$ gives

$$\tilde{e}_i \bar{\pi}_\lambda(b) = \tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda + qL(\lambda) \in B(\lambda) \cup \{0\}.$$

Therefore, by Lemma 7.19 and Corollary 7.14(c), we conclude $\tilde{e}_i b = \tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1 \in B(\infty) \cup \{0\}$. $\square$

Proposition 7.23 ($\mathbf{O}(r)$). *Let $\lambda \in P^+$ and $\alpha \in Q_+(r)$. If $b \in B(\infty)_{-\alpha}$ satisfies $\bar{\pi}_\lambda(b) \neq 0$, then we have $\tilde{e}_i \bar{\pi}_\lambda(b) = \bar{\pi}_\lambda(\tilde{e}_i b)$.*

Proof. Let $P = \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1 \in L(\infty)_{-\alpha}$, $b = \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1 + qL(\infty) \in B(\infty)_{-\alpha}$ and $u = \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda$. Then by Proposition 7.12, we have $u \equiv Pv_\lambda \pmod{qL(\lambda)}$.

Since $\bar{\pi}_\lambda(b) \neq 0$, we have $u \notin qL(\lambda)$. Observe that for any $\mu \in P^+$, Lemma 7.5(g) gives

$$\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} (v_\lambda \otimes v_\mu) \equiv u \otimes v_\mu \pmod{qL(\lambda) \otimes L(\mu)}.$$

Hence by Lemma 7.21,

$$(7.9) \quad \tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} (v_\lambda \otimes v_\mu) \equiv \tilde{e}_i u \otimes v_\mu \pmod{qL(\lambda) \otimes L(\mu)}.$$

On the other hand, for any $\mu \gg 0$, Lemma 7.19 yields

$$\tilde{e}_i (\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_{\lambda+\mu}) \equiv \tilde{e}_i (Pv_{\lambda+\mu}) \equiv (\tilde{e}_i P)v_{\lambda+\mu} \pmod{qL(\lambda+\mu)}.$$

Applying $\Phi_{\lambda, \mu}$, Proposition 7.6 yields

$$\tilde{e}_i (\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} (v_\lambda \otimes v_\mu)) \equiv (\tilde{e}_i P)(v_\lambda \otimes v_\mu) \pmod{qL(\lambda) \otimes L(\mu)}.$$

Together with (7.9), we obtain

$$\tilde{e}_i u \otimes v_\mu \equiv (\tilde{e}_i P)(v_\lambda \otimes v_\mu) \pmod{qL(\lambda) \otimes L(\mu)}.$$

Therefore, by applying $S_{\lambda, \mu}$, we conclude $\tilde{e}_i u \equiv (\tilde{e}_i P)v_\lambda$, which gives

$$\tilde{e}_i \bar{\pi}_\lambda(b) = \bar{\pi}_\lambda(\tilde{e}_i b),$$

as claimed. $\square$

Proposition 7.24 (**(B(r))**). *For every $\lambda \in P^+$ and $\alpha \in Q_+(r)$, we have*

$$\tilde{e}_i B(\lambda)_{\lambda-\alpha} \subset B(\lambda) \cup \{0\}.$$

Proof. Our assertion is an immediate consequence of Proposition 7.23, Corollary 7.14(b) and Proposition 7.22. $\square$

To prove the remaining inductive statements, we use the symmetric form on $V(\lambda)$. Since the situation is special when $a_{ii} = 0$, we shall introduce the operator Q_i on $V(\lambda)$ by

$$(7.10) \quad Q_i(f_i^{(n)} u) = \begin{cases} (n+1)f_i^{(n)} u & \text{if } a_{ii} = 0, \\ f_i^{(n)} u & \text{otherwise} \end{cases}$$

for $u \in \text{Ker } e_i$ so that $Q_i = \text{id}_{V(\lambda)}$ when $a_{ii} \neq 0$.

Lemma 7.25. *Let $\lambda \in P^+$ be a dominant integral weight and let $(\ , \)$ be the symmetric bilinear form on $V(\lambda)$ given by (2.3). Then we have*

$$(L(\lambda)_{\lambda-\alpha}, L(\lambda)_{\lambda-\alpha}) \subset \mathbf{A}_0$$

for every $\alpha \in Q_+(r)$. Moreover, we have

$$(7.11) \quad (\tilde{f}_i u, v) \equiv (u, Q_i \tilde{e}_i v) \pmod{q\mathbf{A}_0}$$

for every $\alpha \in Q_+(r)$, $u \in L(\lambda)_{\lambda-\alpha+\alpha_i}$ and $v \in L(\lambda)_{\lambda-\alpha}$.

Proof. We shall argue by the induction on $|\alpha|$. By Corollary 7.20 (a), it is enough to show (7.11) for $u = f_i^{(n)} u_0 \neq 0$, $v = f_i^{(m)} v_0 \neq 0$ with $u_0 \in \text{Ker } e_i \cap L(\lambda)$ and $v_0 \in \text{Ker } e_i \cap L(\mu)$.

Set $a = \langle h_i, \text{wt}(v_0) \rangle$. If $a = 0$ or $m = 0$, then the assertion is trivial. Assume that $a > 0$ and $m > 0$. Then we have

$$(\tilde{f}_i u, v) = (f_i^{(n+1)} u_0, f_i^{(m)} v_0) = a(f_i^{(n)} u_0, q_i^{-1} K_i e_i f_i^{(m)} v_0).$$

Here, $a = [n+1]_i^{-1}$ or 1 according as $a_{ii} = 2$ or $a_{ii} \leq 0$. By (4.6), we have

$$a q_i^{-1} K_i e_i f_i^{(m)} v_0 \in \begin{cases} (1 + q \mathbf{A}_0) f_i^{(m-1)} v_0 & \text{if } a_{ii} \neq 0, \\ m(1 + q \mathbf{A}_0) f_i^{(m-1)} v_0 & \text{if } a_{ii} = 0. \end{cases}$$

Hence we have $(\tilde{f}_i u, v) \in (1 + q \mathbf{A}_0)(u, Q_i \tilde{e}_i v)$. By the induction hypothesis, we have $(u, Q_i \tilde{e}_i v) \in \mathbf{A}_0$, and we are done. $\square$

Lemma 7.26. *Let $\alpha = \sum_{k=1}^r \alpha_{i_k} \in Q_+$, $P, Q \in U_q^-(\mathfrak{g})_{-\alpha}$ and $m \in \mathbf{Z}$. Then for $\lambda \gg 0$, we have*

$$(P, Q) \equiv \prod_{k=1}^r (1 - q_{i_k}^2)^{-1} (P v_\lambda, Q v_\lambda) \pmod{q^m \mathbf{A}_0}.$$

Proof. If $P = 1$, our assertion is clear. We shall argue by the induction on the height r of α . We may assume $P = f_i R$. Then we have

$$\begin{aligned} (f_i R v_\lambda, Q v_\lambda) &= (R v_\lambda, q_i^{-1} K_i e_i Q v_\lambda) \\ &= \left(R v_\lambda, q_i^{-1} K_i \left(Q e_i + \frac{K_i^{-1}(e'_i Q) - K_i(e''_i Q)}{q_i^{-1} - q_i} \right) v_\lambda \right) \\ &= \left(R v_\lambda, \frac{(e'_i Q) v_\lambda - K_i^2(e''_i Q) v_\lambda}{1 - q_i^2} \right) \\ &= \left(R v_\lambda, \frac{(e'_i Q) v_\lambda - q_i^{2\langle h_i, \lambda - \alpha + \alpha_i \rangle} (e''_i Q) v_\lambda}{1 - q_i^2} \right) \\ &= (1 - q_i^2)^{-1} (R v_\lambda, (e'_i Q) v_\lambda) - q_i^{2\langle h_i, \lambda - \alpha + \alpha_i \rangle} (1 - q_i^2)^{-1} (R v_\lambda, (e''_i Q) v_\lambda), \\ &\equiv (1 - q_i^2)^{-1} (R, e'_i Q) = (1 - q_i^2)^{-1} (f_i R, Q) \pmod{q^m \mathbf{A}_0} \end{aligned}$$

by the induction hypothesis. $\square$

For a finitely generated $\mathbf{A}_0$ -submodule L of $V(\lambda)_{\lambda-\alpha}$, we set

$$L^\vee := \{u \in V(\lambda)_{\lambda-\alpha}; (u, L) \subset \mathbf{A}_0\}.$$

Then we have $(L^\vee)^\vee = L$. Similarly, we define $L^\vee$ for a finitely generated $\mathbf{A}_0$ -submodule L of $U_q^-(\mathfrak{g})_{-\alpha}$.

Lemma 7.27. *If $\lambda \gg 0$, then for any $\alpha \in Q_+(r)$ we have $\pi_\lambda(L(\infty)_{-\alpha}^\vee) = L(\lambda)_{\lambda-\alpha}^\vee$.*

Proof. Let $\{P_k\}$ be an $\mathbf{A}_0$ -basis of $L(\infty)_{-\alpha}$, and $\{Q_k\}$ the dual basis: $(P_k, Q_j) = \delta_{kj}$. Then $L(\infty)_{-\alpha}^\vee = \sum_j \mathbf{A}_0 Q_j$. By Proposition 7.13, we have $L(\lambda)_{\lambda-\alpha} = \sum_k \mathbf{A}_0 P_k v_\lambda$. By Lemma 7.26, we have $(P_k v_\lambda, Q_j v_\lambda) \equiv \delta_{kj} \pmod{q\mathbf{A}_0}$ for $\lambda \gg 0$. Hence we conclude

$$L(\lambda)_{\lambda-\alpha}^\vee = \sum_j \mathbf{A}_0 Q_j v_\lambda = \pi_\lambda(L(\infty)_{-\alpha}^\vee) \quad \text{for } \lambda \gg 0.$$

□

In the following lemma, we will prove a special case of $\mathbf{F}(r)$.

Lemma 7.28. *Let $\mu \gg 0$ and $\alpha \in Q_+(r)$. If $\lambda \in P^+$, then we have*

$$\Psi_{\lambda,\mu}((L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha}) \subset L(\lambda+\mu)_{\lambda+\mu-\alpha}.$$

Proof. Recall that Lemma 7.8 gives

$$(L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha} = \sum_i \tilde{f}_i((L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha+\alpha_i}) + v_\lambda \otimes L(\mu)_{\mu-\alpha}.$$

By the induction hypothesis $\mathbf{F}(r-1)$, we have

$$\begin{aligned} \Psi_{\lambda,\mu}(\sum_i \tilde{f}_i(L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha+\alpha_i}) &= \sum_i \tilde{f}_i \Psi_{\lambda,\mu}((L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha+\alpha_i}) \\ &\subset \sum_i \tilde{f}_i L(\lambda+\mu)_{\lambda+\mu-\alpha+\alpha_i} = L(\lambda+\mu)_{\lambda+\mu-\alpha}. \end{aligned}$$

Thus it remains to show

$$\Psi_{\lambda,\mu}(v_\lambda \otimes L(\mu)_{\mu-\alpha}) \subset L(\lambda+\mu)_{\lambda+\mu-\alpha}.$$

For $u \in L(\lambda+\mu)_{\lambda+\mu-\alpha}^\vee$, by Lemma 7.27, we can write $u = P v_{\lambda+\mu}$ for some $P \in L(\infty)_{-\alpha}^\vee$. Observe that

$$\Delta(P) = P \otimes 1 + (\text{intermediate terms}) + K_\alpha \otimes P,$$

which yields

$$\begin{aligned}
(\Phi_{\lambda,\mu}(u), v_\lambda \otimes L(\mu)_{\mu-\alpha}) &= (\Delta(P)(v_\lambda \otimes v_\mu), v_\lambda \otimes L(\mu)_{\mu-\alpha}) \\
&= (Pv_\lambda \otimes v_\mu + \cdots + K_\alpha v_\lambda \otimes Pv_\mu, v_\lambda \otimes L(\mu)_{\mu-\alpha}) \\
&= q^{(\alpha|\lambda)}(Pv_\mu, L(\mu)_{\mu-\alpha}).
\end{aligned}$$

Since $\mu \gg 0$, Lemma 7.27 implies that $Pv_\mu \in L(\mu)^\vee$. Thus we have

$$\begin{aligned}
(u, \Psi_{\lambda,\mu}(v_\lambda \otimes L(\mu)_{\mu-\alpha})) &= (\Phi_{\lambda,\mu}(u), v_\lambda \otimes L(\mu)_{\mu-\alpha}) \\
&= q^{(\alpha|\lambda)}(Pv_\mu, L(\mu)_{\mu-\alpha}) \subset \mathbf{A}_0.
\end{aligned}$$

Hence, we have

$$\Psi_{\lambda,\mu}(v_\lambda \otimes L(\mu)_{\mu-\alpha}) \subset L(\lambda + \mu)_{\lambda+\mu-\alpha}^{\vee\vee} = L(\lambda + \mu)_{\lambda+\mu-\alpha},$$

which completes the proof. $\square$

Proposition 7.29 (**J**(r)). *Suppose $\alpha \in Q_+(r)$ and $b \in B(\infty)_{-\alpha}$. If $\tilde{e}_i b \neq 0$, then we have $b = \tilde{f}_i \tilde{e}_i b$.*

Proof. Let $b = \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1 \in B(\infty)_{-\alpha}$ and suppose $\tilde{e}_i b = b' \neq 0$. If $i_1 = \cdots = i_r$, then $\tilde{e}_i b' \neq 0$ implies that $i_1 = \cdots = i_r = i$, and our assertion follows immediately.

If there exists a positive integer $t < r$ such that $i_t \neq i_{t+1} = \cdots = i_r$, take $\mu \gg 0$ and set $\lambda_0 = \Lambda_{i_t}$, $\lambda = \lambda_0 + \mu$. Then Lemma 7.7 yields

$$\tilde{f}_{i_1} \cdots \tilde{f}_{i_r}(v_{\lambda_0} \otimes v_\mu) \equiv v \otimes v' \pmod{qL(\lambda_0) \otimes L(\mu)}$$

for some $v \in L(\lambda_0)_{\lambda_0-\beta}$ and $v' \in L(\mu)_{\mu-\gamma}$ and $\beta, \gamma \in Q_+(r-1) \setminus \{0\}$ with $\alpha = \beta + \gamma$ such that $v + qL(\lambda_0) \in B(\lambda_0) \cup \{0\}$ and $v' + qL(\mu) \in B(\mu) \cup \{0\}$. By Corollary 7.20 (b), we have

$$\tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r}(v_{\lambda_0} \otimes v_\mu) \equiv \tilde{e}_i(v \otimes v') \pmod{qL(\lambda_0) \otimes L(\mu)}.$$

Applying $\Psi_{\lambda_0,\mu}$, **G**($r-1$) yields

$$\pi_\lambda(\tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} \cdot 1) = \tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_{\lambda_0+\mu} \equiv \Psi_{\lambda_0,\mu}(\tilde{e}_i(v \otimes v')) \pmod{qL(\lambda)}.$$

Thus $\tilde{e}_i(v \otimes v') \notin qL(\lambda_0) \otimes L(\mu)$, and Lemma 7.5 (d) gives

$$\begin{aligned} \tilde{f}_{i_1} \cdots \tilde{f}_{i_r}(v_{\lambda_0} \otimes v_\mu) &\equiv v \otimes v' \equiv \tilde{f}_i \tilde{e}_i(v \otimes v') \\ &\equiv \tilde{f}_i \tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r}(v_{\lambda_0} \otimes v_\mu) \pmod{qL(\lambda_0) \otimes L(\mu)}. \end{aligned}$$

Hence, by applying $\Psi_{\lambda_0, \mu}$, Lemma 7.28 gives

$$\tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda \equiv \tilde{f}_i \tilde{e}_i \tilde{f}_{i_1} \cdots \tilde{f}_{i_r} v_\lambda \pmod{qL(\lambda)}.$$

Therefore by Lemma 7.11 and Corollary 7.14 (c), we have $b \equiv \tilde{f}_i \tilde{e}_i b \pmod{qL(\infty)}$. $\square$

Proposition 7.30 (C(r)). *Suppose $\alpha \in Q_+(r)$ and $\lambda \in P^+$. Then for every $i \in I$, $b \in B(\lambda)_{\lambda-\alpha+\alpha_i}$ and $b' \in B(\lambda)_{\lambda-\alpha}$, we have $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$.*

Proof. ($\implies$) Let $\tilde{f}_i b = b' \in B(\lambda)_{\lambda-\alpha}$. Since $b \in B(\lambda)_{\lambda-\alpha+\alpha_i}$, by Lemma 7.3(b), there exists $n \geq 0$ such that $b \equiv f_i^{(n)} u$ with $e_i u = 0$. Hence $b' \equiv f_i^{(n+1)} u$ and $\tilde{e}_i b' \equiv f_i^{(n)} u \equiv b$.

($\impliedby$) Let $b' \in B(\lambda)_{\lambda-\alpha}$ and $b = \tilde{e}_i b' \in B(\lambda)$. Then we have $b' = \pi_\lambda(b'')$ for some $b'' \in L(\infty)_{-\alpha}$ by Corollary 7.14(b). By Proposition 7.23, we have

$$\pi_\lambda(\tilde{e}_i b'') = \tilde{e}_i \pi_\lambda(b'') = \tilde{e}_i b' \neq 0.$$

Hence we have $\tilde{e}_i b'' \neq 0$ and Proposition 7.29 yields $b'' = \tilde{f}_i \tilde{e}_i b''$. By applying π_λ , we obtain

$$\tilde{f}_i \tilde{e}_i b' = \tilde{f}_i \pi_\lambda(\tilde{e}_i b'') = \pi_\lambda(\tilde{f}_i \tilde{e}_i b'') = \pi_\lambda(b'') = b',$$

as claimed. $\square$

Proposition 7.31 (D(r), K(r)). *Let $\alpha \in Q_+(r)$.*

- (a) *For $\lambda \in P^+$, $B(\lambda)_{\lambda-\alpha}$ is a $\mathbf{Q}$ -basis of $L(\lambda)_{\lambda-\alpha}/qL(\lambda)_{\lambda-\alpha}$.*
- (b) *$B(\infty)_{-\alpha}$ is a $\mathbf{Q}$ -basis of $L(\infty)_{-\alpha}/qL(\infty)_{-\alpha}$.*

Proof. We will prove (a) only. The proof of (b) is similar.

Suppose that we have a $\mathbf{Q}$ -linear dependence relation

$$\sum_{b \in B(\lambda)_{\lambda-\alpha}} a_b b = 0 \text{ with } a_b \in \mathbf{Q}.$$

Since $\tilde{e}_i B(\lambda)_{\lambda-\alpha} \subset B(\lambda) \cup \{0\}$ for every $i \in I$, we get

$$0 = \tilde{e}_i \left(\sum_{b \in B(\lambda)_{\lambda-\alpha}} a_b b \right) = \sum_{\substack{b \in B(\lambda)_{\lambda-\alpha}, \\ \tilde{e}_i b \neq 0}} a_b (\tilde{e}_i b).$$

By **D**($r-1$) and **C**(r), the family $\{\tilde{e}_i b; b \in B(\lambda)_{\lambda-\alpha}, \tilde{e}_i b \neq 0\}$ is linearly independent over $\mathbf{Q}$. Therefore $a_b = 0$ whenever $\tilde{e}_i b \neq 0$.

But for each $b \in B(\lambda)_{\lambda-\alpha}$, there exists $i \in I$ such that $\tilde{e}_i b \neq 0$. Hence $a_b = 0$ for every $b \in B(\lambda)_{\lambda-\alpha}$, which completes the proof. $\square$

Lemma 7.32. *Let $\lambda \in P^+$ and $\alpha \in Q_+(r) \setminus \{0\}$.*

- (a) *If $u \in L(\lambda)_{\lambda-\alpha}/qL(\lambda)_{\lambda-\alpha}$ satisfies $\tilde{e}_i u = 0$ for every $i \in I$, then $u = 0$.*
- (b) *If $u \in V(\lambda)_{\lambda-\alpha}$ satisfies $\tilde{e}_i u \in L(\lambda)$ for every $i \in I$, then $u \in L(\lambda)$.*
- (c) *If $u \in L(\infty)_{-\alpha}/qL(\infty)_{-\alpha}$ satisfies $\tilde{e}_i u = 0$ for every $i \in I$, then $u = 0$.*
- (d) *If $u \in U_q^-(\mathfrak{g})_{-\alpha}$ satisfies $\tilde{e}_i u \in L(\infty)$ for every $i \in I$, then $u \in L(\infty)$.*

Proof. (a) Write $u = \sum_{b \in B(\lambda)_{\lambda-\alpha}} a_b b$ with $a_b \in \mathbf{Q}$. Then for every $i \in I$, we have

$$\sum_{\tilde{e}_i b \neq 0} a_b (\tilde{e}_i b) = 0.$$

By the same argument as in the proof of Proposition 7.31, all a_b vanish, which implies $u = 0$.

(b) Choose the smallest $N \geq 0$ such that $q^N u \in L(\lambda)$. If $N > 0$, then $\tilde{e}_i(q^N u) = q^N \tilde{e}_i u \in qL(\lambda)$ for every $i \in I$. Hence, by (a), we would have $q^N u \in qL(\lambda)$; that is, $q^{N-1} u \in L(\lambda)$, a contradiction to the minimality of N . Therefore $N = 0$ and $u \in L(\lambda)$.

The proofs of (c) and (d) are similar. $\square$

Proposition 7.33 (**F**(r)). *For every $\lambda, \mu \in P^+$, we have*

$$\Psi_{\lambda, \mu}((L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha}) \subset L(\lambda + \mu).$$

Proof. We may assume that $\alpha \neq 0$. By Corollary 7.20(b), we have

$$\tilde{e}_i((L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha}) \subset (L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha+\alpha_i}$$

for every $i \in I$. Then by applying $\Psi_{\lambda,\mu}$, the statement $\mathbf{F}(r-1)$ yields

$$\tilde{e}_i \Psi_{\lambda,\mu}((L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha}) \subset \Psi_{\lambda,\mu}((L(\lambda) \otimes L(\mu))_{\lambda+\mu-\alpha+\alpha_i}) \subset L(\lambda+\mu)$$

for every $i \in I$. Hence our assertion follows from Lemma 7.32(b). $\square$

Proposition 7.34 ($\mathbf{N}(r)$). *For all $\alpha \in Q_+(r)$ and $\lambda \in P^+$, we have*

$$\{b \in B(\infty)_{-\alpha}; \bar{\pi}_\lambda(b) \neq 0\} \xrightarrow{\sim} B(\lambda)_{\lambda-\alpha}.$$

Proof. We may assume $\alpha \neq 0$. We already know $\bar{\pi}_\lambda B(\infty)_{-\alpha} \setminus \{0\} = B(\lambda)_{\lambda-\alpha}$. Hence it remains to prove that, for $b, b' \in B(\infty)_{-\alpha}$, $\bar{\pi}_\lambda(b) = \bar{\pi}_\lambda(b') \neq 0$ implies $b = b'$. Choose $i \in I$ such that $\tilde{e}_i \bar{\pi}_\lambda b \neq 0$. By Proposition 7.23, we have

$$\bar{\pi}_\lambda(\tilde{e}_i b) = \bar{\pi}_\lambda(\tilde{e}_i b') \neq 0.$$

Then by the induction hypothesis $\mathbf{N}(r-1)$ and Proposition 7.24, we have $\tilde{e}_i b = \tilde{e}_i b' \neq 0$. Hence Proposition 7.29 yields $b = b'$. $\square$

So far, we have proved all the statements except $\mathbf{G}(r)$. Using these statements, we can prove that Lemma 7.5 holds for all $\alpha \in Q_+(r)$. In particular, we have

Lemma 7.35.

(a) *For all $i \in I$, $\lambda, \mu \in P^+$ and $\alpha \in Q_+(r)$, we have*

$$\tilde{e}_i(B(\lambda) \otimes B(\mu))_{\mu-\alpha} \subset (B(\lambda) \otimes B(\mu)) \cup \{0\}.$$

(b) *If $b \otimes b' \in (B(\lambda) \otimes B(\mu))_{\mu-\alpha}$ and $\tilde{e}_i(b \otimes b') \neq 0$, then*

$$b \otimes b' \equiv \tilde{f}_i \tilde{e}_i(b \otimes b').$$

Finally, we complete the grand-loop argument by proving the statement $\mathbf{G}(r)$.

Proposition 7.36 ($\mathbf{G}(r)$). *For all $i \in I$, $\lambda, \mu \in P^+$ and $\alpha \in Q_+(r)$, we have*

$$\Psi_{\lambda,\mu}((B(\lambda) \otimes B(\mu))_{\lambda+\mu-\alpha}) \subset B(\lambda+\mu) \cup \{0\}.$$

Proof. We may assume $\alpha \neq 0$. Let $b \otimes b' \in (B(\lambda) \otimes B(\mu))_{\lambda+\mu-\alpha}$. If $\tilde{e}_i(b \otimes b') = 0$ for every $i \in I$, then $\tilde{e}_i \Psi_{\lambda,\mu}(b \otimes b') = 0$ for every $i \in I$, and Lemma 7.32 implies $\Psi_{\lambda,\mu}(b \otimes b') = 0$.

If $\tilde{e}_i(b \otimes b') \neq 0$ for some $i \in I$, then by Lemma 7.35 and $\mathbf{G}(r-1)$, we have

$$\begin{aligned} \Psi_{\lambda, \mu}(b \otimes b') &\equiv \Psi_{\lambda, \mu}(\tilde{f}_i \tilde{e}_i(b \otimes b')) \equiv \tilde{f}_i \Psi_{\lambda, \mu}(\tilde{e}_i(b \otimes b')) \\ &\in \tilde{f}_i(B(\lambda + \mu) \cup \{0\}) \subset B(\lambda + \mu) \cup \{0\}. \end{aligned}$$

□

Thus the proof of Theorem 7.1 is completed.

Let $(\ , \)_0$ be the $\mathbf{Q}$ -valued symmetric bilinear form on $L(\lambda)/qL(\lambda)$ given by taking the crystal limit of $(\ , \)$ on $L(\lambda)$. Then we have:

Corollary 7.37. *Let $V(\lambda)$ be the irreducible highest weight $U_q(\mathfrak{g})$ -module with highest weight $\lambda \in P^+$ and let $(L(\lambda), B(\lambda))$ be the crystal basis of $V(\lambda)$.*

(a) *The crystal $B(\lambda)$ forms an orthogonal basis of $L(\lambda)/qL(\lambda)$ with respect to $(\ , \)_0$, and $(b, b)_0 \in \mathbf{Z}_{>0}$ for every $b \in B(\lambda)$. In particular, $(\ , \)_0$ is positive definite on $L(\lambda)/qL(\lambda)$.*

(b) *We have*

$$\begin{aligned} L(\lambda) &= \{u \in V(\lambda); (u, L(\lambda)) \subset \mathbf{A}_0\} \\ &= \{u \in V(\lambda); (u, u) \in \mathbf{A}_0\}. \end{aligned}$$

Proof. (a) We will prove $(b, b')_0 \in \delta_{b, b'} \mathbf{Z}_{>0}$ for all $b, b' \in B(\lambda)_{\lambda - \alpha}$, $\alpha \in Q_+(r)$ by induction on $|\alpha|$. If $|\alpha| = 0$, our assertion is trivial.

If $|\alpha| > 0$, choose $i \in I$ such that $\tilde{e}_i b \neq 0$. Then by Theorem 7.1 and Proposition 7.30, we have

$$(b, b')_0 = (\tilde{f}_i \tilde{e}_i b, b')_0 = (\tilde{e}_i b, Q_i \tilde{e}_i b')_0 \in \mathbf{Z}_{>0}(\tilde{e}_i b, \tilde{e}_i b')_0 \subset \delta_{\tilde{e}_i b, \tilde{e}_i b'} \mathbf{Z}_{>0} = \delta_{b, b'} \mathbf{Z}_{>0},$$

which completes the proof.

(b) Let

$$\begin{aligned} L_1 &= \{u \in V(\lambda); (u, L(\lambda)) \subset \mathbf{A}_0\}, \\ L_2 &= \{u \in V(\lambda); (u, u) \in \mathbf{A}_0\}. \end{aligned}$$

It is clear that $L(\lambda) \subset L_1$ and $L(\lambda) \subset L_2$. Let $u \in V(\lambda)$ be such that $(u, L(\lambda)) \subset \mathbf{A}_0$. Take the smallest $n \geq 0$ such that $q^n u \in L(\lambda)$. If $n > 0$, then $(q^n u, L(\lambda)) \equiv 0$

mod $q\mathbf{A}_0$. Since $(\ , \)_0$ is non-degenerate, $q^n u \equiv 0 \pmod{qL(\lambda)}$; that is $q^{n-1}u \in L(\lambda)$, which contradicts the minimality of n . Hence $n = 0$ and $u \in L(\lambda)$, which proves $L_1 \subset L(\lambda)$.

Similarly, let $u \in V(\lambda)$ be such that $(u, u) \in \mathbf{A}_0$. Take the smallest $n \geq 0$ such that $q^n u \in L(\lambda)$. If $n > 0$, then $(q^n u, q^n u) = q^{2n}(u, u) \in q\mathbf{A}_0$. Since $(\ , \)_0$ is positive definite, we obtain $q^n u \equiv 0 \pmod{qL(\lambda)}$, a contradiction. Hence $n = 0$ and $u \in L(\lambda)$, which proves $L_2 \subset L(\lambda)$. $\square$

Remark 7.38. If $a_{ii} \neq 0$ for every $i \in I$, then $B(\lambda)$ is an orthonormal basis of $L(\lambda)/qL(\lambda)$.

The following lemma immediately follows from the preceding proposition and Lemma 7.26.

Lemma 7.39.

- (a) $(L(\infty), L(\infty)) \subset \mathbf{A}_0$.
- (b) Let $(\ , \)_0$ be the $\mathbf{Q}$ -valued inner product on $L(\infty)/qL(\infty)$ induced by taking the crystal limit of $(\ , \)$ on $L(\infty)$. Then $B(\infty)$ is an orthogonal basis of $L(\infty)/qL(\infty)$. In particular, $(\ , \)_0$ is positive definite on $L(\infty)/qL(\infty)$.
- (c) We have

$$\begin{aligned} L(\infty) &= \{P \in U_q^-(\mathfrak{g}); (u, L(\infty)) \subset \mathbf{A}_0\} \\ &= \{u \in U_q^-(\mathfrak{g}); (u, u) \in \mathbf{A}_0\}. \end{aligned}$$

Corollary 7.40. For every $P \in L(\infty)$, we have $P^* \in L(\infty)$.

Proof. Let $P \in L(\infty)$. Then by Lemma 7.39 (a) and (6.6), we have $(P^*, P^*) = (P, P) \in \mathbf{A}_0$. Therefore, by Lemma 7.39 (c), we have $P^* \in L(\infty)$. $\square$

8. Balanced Triple

In the following three sections, we will *globalize* the theory of crystal bases. Recall that $\mathbf{A}_0$ is the subring of $\mathbf{Q}(q)$ consisting of rational functions in q that are regular

at $q = 0$. Similarly, let $\mathbf{A}_\infty$ be the subring of $\mathbf{Q}(q)$ consisting of rational functions in q that are regular at $q = \infty$. Thus we have

$$\mathbf{A}_0/q\mathbf{A}_0 \xrightarrow{\sim} \mathbf{Q} \quad \text{and} \quad \mathbf{A}_\infty/q^{-1}\mathbf{A}_\infty \xrightarrow{\sim} \mathbf{Q}$$

by evaluation at $q = 0$ and $q = \infty$, respectively. Finally, we denote by $\mathbf{A} = \mathbf{Q}[q, q^{-1}]$, the ring of Laurent polynomials in q .

For a subring A of $\mathbf{Q}(q)$ and a vector space V over $\mathbf{Q}(q)$, an A -lattice is, by definition, a free A -module of V generating V as a vector space over $\mathbf{Q}(q)$. Assume that

$$(8.1) \quad \begin{aligned} &V \text{ is a vector space over } \mathbf{Q}(q), \text{ and } V^{\mathbf{A}}, L_0 \text{ and } L_\infty \text{ are an } \mathbf{A}\text{-lattice,} \\ &\text{an } \mathbf{A}_0\text{-lattice and an } \mathbf{A}_\infty\text{-lattice of } V, \text{ respectively.} \end{aligned}$$

Note that $(V^{\mathbf{A}} \cap L_0)/(V^{\mathbf{A}} \cap qL_0) \rightarrow L_0/qL_0$ is an isomorphism.

Definition 8.1. Set $E = V^{\mathbf{A}} \cap L_0 \cap L_\infty$. Then a triple $(V^{\mathbf{A}}, L_0, L_\infty)$ is called a *balanced triple* for V if the canonical homomorphisms $\mathbf{A} \otimes E \rightarrow V^{\mathbf{A}}$, $\mathbf{A}_0 \otimes E \rightarrow L_0$ and $\mathbf{A}_\infty \otimes E \rightarrow L_\infty$ are isomorphisms.

Theorem 8.2. Let $V, V^{\mathbf{A}}, L_0$ and L_∞ be as in (8.1). Set $E = V^{\mathbf{A}} \cap L_0 \cap L_\infty$. Then the following statements are equivalent.

- (a) $(V^{\mathbf{A}}, L_0, L_\infty)$ is a balanced triple.
- (b) The canonical map $E \rightarrow L_0/qL_0$ is an isomorphism.
- (c) The canonical map $E \rightarrow L_\infty/q^{-1}L_\infty$ is an isomorphism.

Let $(V^{\mathbf{A}}, L_0, L_\infty)$ be a balanced triple for a $\mathbf{Q}(q)$ -vector space V , and let

$$G: L_0/qL_0 \longrightarrow E := V^{\mathbf{A}} \cap L_0 \cap L_\infty$$

be the inverse of the canonical isomorphism $E \xrightarrow{\sim} L_0/qL_0$. If B is a $\mathbf{Q}$ -basis of L_0/qL_0 , then $G(B) := \{G(b); b \in B\}$ is a $\mathbf{Q}$ -basis of E . Therefore $G(B)$ is a $\mathbf{Q}(q)$ -basis of V , an $\mathbf{A}$ -basis of $V^{\mathbf{A}}$, an $\mathbf{A}_0$ -basis of L_0 and an $\mathbf{A}_\infty$ -basis of L_∞ as well.

Definition 8.3. We call $G(B)$ the *global basis* of V corresponding to the *local basis* B of V at $q = 0$.

Our goal is to construct the global basis $G(\lambda)$ of $V(\lambda)$ (resp. $G(\infty)$ of $U_q^-(\mathfrak{g})$) corresponding to the local basis $B(\lambda)$ of $V(\lambda)$ (resp. $B(\infty)$ of $U_q^-(\mathfrak{g})$) at $q = 0$. For this purpose, we need the following lemma which is proved in [10].

Lemma 8.4 ([10]). *Let V , $V^{\mathbf{A}}$, L_0 and L_∞ be as in (8.1). Let F be a $\mathbf{Q}$ -vector subspace of $V^{\mathbf{A}} \cap L_0 \cap L_\infty$ satisfying the following conditions:*

- (a) *the canonical maps $F \longrightarrow L_0/qL_0$ and $F \longrightarrow L_\infty/q^{-1}L_\infty$ are injective.*
- (b) *$V^{\mathbf{A}} = \mathbf{A}F$.*

Then $(V^{\mathbf{A}}, L_0, L_\infty)$ is a balanced triple and $F = V^{\mathbf{A}} \cap L_0 \cap L_\infty$.

9. Global Bases

Let $U_{\mathbf{A}}^0(\mathfrak{g})$ be the $\mathbf{A}$ -subalgebra of $U_q(\mathfrak{g})$ generated by q^h , $\prod_{k=1}^m \frac{1 - q^k \cdot q^h}{1 - q^k}$ ($h \in P^\vee$, $m \in \mathbf{Z}_{>0}$). We denote by $U_{\mathbf{A}}^+(\mathfrak{g})$ (resp. $U_{\mathbf{A}}^-(\mathfrak{g})$) the $\mathbf{A}$ -subalgebra of $U_q(\mathfrak{g})$ generated by $e_i^{(n)}$ (resp. $f_i^{(n)}$) for $i \in I$, $n \in \mathbf{Z}_{\geq 0}$. Let $U_{\mathbf{A}}(\mathfrak{g})$ be the $\mathbf{A}$ -subalgebra of $U_q(\mathfrak{g})$ generated by $U_{\mathbf{A}}^0(\mathfrak{g})$, $U_{\mathbf{A}}^+(\mathfrak{g})$ and $U_{\mathbf{A}}^-(\mathfrak{g})$.

Then we can show that

$$(9.1) \quad U_{\mathbf{A}}(\mathfrak{g}) \simeq U_{\mathbf{A}}^-(\mathfrak{g}) \otimes U_{\mathbf{A}}^0(\mathfrak{g}) \otimes U_{\mathbf{A}}^+(\mathfrak{g}).$$

Let $V(\lambda)$ be the irreducible highest weight $U_q(\mathfrak{g})$ -module with highest weight $\lambda \in P^+$ and highest weight vector v_λ , and let

$$V(\lambda)^{\mathbf{A}} = U_{\mathbf{A}}(\mathfrak{g})v_\lambda = U_{\mathbf{A}}^-(\mathfrak{g})v_\lambda.$$

Consider the $\mathbf{Q}$ -algebra automorphism $\bar{\cdot} : U_q(\mathfrak{g}) \rightarrow U_q(\mathfrak{g})$ defined by

$$(9.2) \quad q \mapsto q^{-1}, \quad q^h \mapsto q^{-h}, \quad e_i \mapsto e_i, \quad f_i \mapsto f_i$$

for $h \in P^\vee$ and $i \in I$. Then we get a $\mathbf{Q}$ -linear automorphism $\bar{\cdot}$ on $V(\lambda)$ defined by

$$(9.3) \quad Pv_\lambda \mapsto \bar{P}v_\lambda \quad \text{for } P \in U_q(\mathfrak{g}).$$

Our goal is to prove the following theorem.

Theorem 9.1.

- (a) *$(U_{\mathbf{A}}^-(\mathfrak{g}), L(\infty), L(\infty)^-)$ is a balanced triple for $U_q^-(\mathfrak{g})$.*

(b) $(V(\lambda)^{\mathbf{A}}, L(\lambda), L(\lambda)^-)$ is a balanced triple for $V(\lambda)$.

Once we have Theorem 9.1, the canonical maps

$$\begin{aligned} U_{\mathbf{A}}^-(\mathfrak{g}) \cap L(\infty) \cap L(\infty)^- &\longrightarrow L(\infty)/qL(\infty), \\ V(\lambda)^{\mathbf{A}} \cap L(\lambda) \cap L(\lambda)^- &\longrightarrow L(\lambda)/qL(\lambda) \end{aligned}$$

are isomorphisms. As in Section 8, let

$$\begin{aligned} G_{\infty}: L(\infty)/qL(\infty) &\longrightarrow U_{\mathbf{A}}^-(\mathfrak{g}) \cap L(\infty) \cap L(\infty)^-, \\ G_{\lambda}: L(\lambda)/qL(\lambda) &\longrightarrow V(\lambda)^{\mathbf{A}} \cap L(\lambda) \cap L(\lambda)^- \end{aligned}$$

be the inverses of the isomorphisms above, and set

$$(9.4) \quad G(\infty) = \{G_{\infty}(b); b \in B(\infty)\}, \quad G(\lambda) = \{G_{\lambda}(b); b \in B(\lambda)\}.$$

Then we have:

Theorem 9.2.

- (a) $G(\infty)$ (resp. $G(\lambda)$) is an $\mathbf{A}$ -basis of $U_{\mathbf{A}}^-(\mathfrak{g})$ (resp. $V(\lambda)^{\mathbf{A}}$).
- (b) $G(\infty)$ (resp. $G(\lambda)$) is an $\mathbf{A}_0$ -basis of $L(\infty)$ (resp. $L(\lambda)$).
- (c) $G(\infty)$ (resp. $G(\lambda)$) is a $\mathbf{Q}(q)$ -basis of $U_q^-(\mathfrak{g})$ (resp. $V(\lambda)$).
- (d) $G(\infty)$ (resp. $G(\lambda)$) is a $\mathbf{Q}[q]$ -basis of $U_{\mathbf{A}}^-(\mathfrak{g}) \cap L(\infty)$ (resp. $V(\lambda)^{\mathbf{A}} \cap L(\lambda)$).

Proof. These are immediate consequences of the definition of a balanced triple. $\square$

Moreover, we have

Theorem 9.3.

- (a) For $b \in B(\infty)$, $G_{\infty}(b)$ is a unique element of $U_{\mathbf{A}}^-(\mathfrak{g}) \cap L(\infty)$ such that

$$G_{\infty}(b) \equiv b \pmod{qL(\infty)}, \quad \overline{G_{\infty}(b)} = G_{\infty}(b).$$

- (b) For $b \in B(\lambda)$, $G_{\lambda}(b)$ is a unique element of $V(\lambda)^{\mathbf{A}} \cap L(\lambda)$ such that

$$G_{\lambda}(b) \equiv b \pmod{qL(\lambda)}, \quad \overline{G_{\lambda}(b)} = G_{\lambda}(b).$$

Proof. The proof of (a) being similar, we will prove only (b). By the definition of G_λ , it is clear that $G_\lambda(b) \equiv b \pmod{qL(\lambda)}$. Set

$$v = \frac{G_\lambda(b) - \overline{G_\lambda(b)}}{q - q^{-1}}.$$

Clearly, $G_\lambda(b)$ and $\overline{G_\lambda(b)}$ are contained in $E := V(\lambda)^{\mathbf{A}} \cap L(\lambda) \cap L(\lambda)^-$. Write $G_\lambda(b) = \sum_j c_j(q) P_j v_\lambda$, where $c_j(q) \in \mathbf{A}$ and P_j is a monomial in $f_i^{(n)}$'s ($i \in I$, $n \in \mathbf{Z}_{>0}$). Then we have $\overline{G_\lambda(b)} = \sum_j c_j(q^{-1}) P_j v_\lambda$, from which it follows that

$$v := \frac{G_\lambda(b) - \overline{G_\lambda(b)}}{q - q^{-1}} = \sum_j \frac{c_j(q) - c_j(q^{-1})}{q - q^{-1}} P_j v_\lambda \in V(\lambda)^{\mathbf{A}}.$$

Since $\frac{1}{q - q^{-1}} = \frac{q}{q^2 - 1} \in q\mathbf{A}_0 \cap \mathbf{A}_\infty$, we have $v \in qL(\lambda) \cap L(\lambda)^-$. Thus v is mapped to 0 under the canonical isomorphism $E \rightarrow L(\lambda)/qL(\lambda)$, which implies $v = 0$. Therefore we get $\overline{G_\lambda(b)} = G_\lambda(b)$.

Finally, suppose that $u \in V(\lambda)^{\mathbf{A}} \cap L(\lambda)$ satisfies

$$u \equiv b \pmod{qL(\lambda)} \quad \text{and} \quad \overline{u} = u.$$

Then $u \in E := V(\lambda)^{\mathbf{A}} \cap L(\lambda) \cap L(\lambda)^-$. Since the canonical map $E \rightarrow L(\lambda)/qL(\lambda)$ is an isomorphism, we must have $u = G_\lambda(b)$. $\square$

Definition 9.4. We call $G(\infty)$ (resp. $G(\lambda)$) the *global basis* of $U_q^-(\mathfrak{g})$ (resp. $V(\lambda)$).

10. Existence of Global Bases

In this section, we will prove Theorem 9.1. For $i \in I$ and $n \geq 1$, define

$$(f_i^n U_q^-(\mathfrak{g}))^{\mathbf{A}} = f_i^n U_q^-(\mathfrak{g}) \cap U_{\mathbf{A}}^-(\mathfrak{g}), \quad (f_i^n V(\lambda))^{\mathbf{A}} = (f_i^n U_q^-(\mathfrak{g}))^{\mathbf{A}} V(\lambda).$$

Then $(f_i^n V(\lambda))^{\mathbf{A}}$ is an $\mathbf{A}$ -lattice of $f_i^n V(\lambda)$.

Lemma 10.1. *Fix $i \in I$. Let $u \in U_q^-(\mathfrak{g})$ and consider the i -string decomposition $u = \sum_{l \geq 0} f_i^{(l)} u_l$ with $e'_i u_l = 0$ for every $l \geq 0$. If $u \in U_{\mathbf{A}}^-(\mathfrak{g})$, then $u_l \in U_{\mathbf{A}}^-(\mathfrak{g})$ for every $l \geq 0$.*

Proof. We first claim that

$$(10.1) \quad e_i'^{(n)} U_{\mathbf{A}}^-(\mathfrak{g}) \subset U_{\mathbf{A}}^-(\mathfrak{g}).$$

Indeed, the commutation relation (6.4) implies that

$$\left\{ Q \in U_q^-(\mathfrak{g}) ; e_i'^{(n)} Q \in U_q^-(\mathfrak{g})^{\mathbf{A}} \text{ for every } n \geq 0 \right\}$$

is stable under the left multiplication by any $f_j^{(m)}$ ($j \in I, m \geq 0$).

Hence the projection operator P_i introduced in (6.7) satisfies $P_i U_{\mathbf{A}}^-(\mathfrak{g}) \subset U_{\mathbf{A}}^-(\mathfrak{g})$. Then Proposition 6.1 implies that all u_l belong to $U_{\mathbf{A}}^-(\mathfrak{g})$. $\square$

Then as an immediate consequence of Lemma 10.1, we have:

Lemma 10.2.

- (a) $(f_i^n U_q^-(\mathfrak{g}))^{\mathbf{A}} := f_i^n U_q^-(\mathfrak{g}) \cap U_{\mathbf{A}}^-(\mathfrak{g}) = \sum_{l \geq n} f_i^{(l)} U_{\mathbf{A}}^-(\mathfrak{g})$.
- (b) $(f_i^n V(\lambda))^{\mathbf{A}} := (f_i^n U_q^-(\mathfrak{g}))^{\mathbf{A}} V(\lambda) = \sum_{l \geq n} f_i^{(l)} V(\lambda)^{\mathbf{A}}$.

In order to prove Theorem 9.1, we shall show the following statements by the induction on r .

X(r) : For every $\alpha \in Q_+(r)$, the triple $(U_{\mathbf{A}}^-(\mathfrak{g})_{-\alpha}, L(\infty)_{-\alpha}, L(\infty)_{-\alpha}^-)$ is balanced.

Y(r) : For all $\lambda \in P^+$ and $\alpha \in Q_+(r)$, the triple $(V(\lambda)_{\lambda-\alpha}, L(\lambda)_{\lambda-\alpha}, L(\lambda)_{\lambda-\alpha}^-)$ is balanced.

Z(r) : For every $\alpha \in Q_+(r)$ and $\lambda \in P^+$, let

$$G_{\lambda} : L(\lambda)_{\lambda-\alpha} / qL(\lambda)_{\lambda-\alpha} \longrightarrow V(\lambda)_{\lambda-\alpha}^{\mathbf{A}} \cap L(\lambda) \cap L(\lambda)^-$$

be the inverse of the canonical isomorphism

$$V(\lambda)_{\lambda-\alpha}^{\mathbf{A}} \cap L(\lambda) \cap L(\lambda)^- \xrightarrow{\sim} L(\lambda)_{\lambda-\alpha} / qL(\lambda)_{\lambda-\alpha}$$

guaranteed by **Y**(r). Then for any $b \in B(\lambda)_{\lambda-\alpha} \cap \tilde{f}_i^n B(\lambda)$ with $n \geq 0$, we have $G_{\lambda}(b) \in (f_i^n V(\lambda))^{\mathbf{A}}$.

If $r = 0$, these assertions are obvious. Suppose $r > 0$ and assume that the statements $\mathbf{X}(r-1)$, $\mathbf{Y}(r-1)$ and $\mathbf{Z}(r-1)$ are true. Then we can define

$$\begin{aligned} G_\infty &: L(\infty)_{-\beta}/qL(\infty)_{-\beta} \xrightarrow{\sim} U_{\mathbf{A}}^-(\mathfrak{g})_{-\beta} \cap L(\infty) \cap L(\infty)^-, \\ G_\lambda &: L(\lambda)_{\lambda-\beta}/qL(\lambda)_{\lambda-\beta} \xrightarrow{\sim} V(\lambda)_{\lambda-\beta}^{\mathbf{A}} \cap L(\lambda) \cap L(\lambda)^- \end{aligned}$$

for $\beta \in Q_+(r-1)$. Thus we have:

Lemma 10.3. *For all $\lambda \in P^+$ and $\beta \in Q_+(r-1)$, we have*

- (a) $U_{\mathbf{A}}^-(\mathfrak{g})_{-\beta} = \bigoplus_{b \in B(\infty)_{-\beta}} \mathbf{A}G_\infty(b)$,
 $V(\lambda)_{\lambda-\beta}^{\mathbf{A}} = \bigoplus_{b \in B(\lambda)_{\lambda-\beta}} \mathbf{A}G_\lambda(b)$,
- (b) $\overline{G_\infty(b)} = G_\infty(b)$ for every $b \in B(\infty)_{-\beta}$,
 $\overline{G_\lambda(b)} = G_\lambda(b)$ for every $b \in B(\lambda)_{\lambda-\beta}$,
- (c) $G_\infty(b)v_\lambda = G_\lambda(\pi_\lambda(b))$ for every $b \in B(\infty)_{-\beta}$.

The following proposition will play a crucial role in proving the statement $\mathbf{X}(r)$ and $\mathbf{Y}(r)$.

Proposition 10.4. *Let $i \in I$, $\alpha \in Q_+(r)$. Then for any $n \geq 1$, the triple*

$$\left((f_i^n V(\lambda))_{\lambda-\alpha}^{\mathbf{A}}, f_i^n V(\lambda) \cap L(\lambda)_{\lambda-\alpha}, f_i^n V(\lambda) \cap L(\lambda)_{\lambda-\alpha}^- \right)$$

is balanced.

Remark 10.5. By Proposition 4.9, we have

$$(f_i^n V(\lambda) \cap L(\lambda)_{\lambda-\alpha}) / (f_i^n V(\lambda) \cap qL(\lambda)_{\lambda-\alpha}) \simeq \bigoplus_{b \in B(\lambda)_{\lambda-\alpha} \cap \tilde{f}_i^n B(\lambda)} \mathbf{Q}b.$$

Proof. If $a_{ii} = 2$, our assertion can be proved as in [10]. Hence we assume that $a_{ii} \leq 0$. In this case, we have

$$(f_i^n V(\lambda))_{\lambda-\alpha}^{\mathbf{A}} = \sum_{l \geq n} f_i^l (V(\lambda)_{\lambda-\alpha+l\alpha_i}^{\mathbf{A}}) = f_i^n (V(\lambda)_{\lambda-\alpha+n\alpha_i}^{\mathbf{A}}).$$

We will use the descending induction on n . Note that if $n > r$, then $(f_i^n V(\lambda))_{\lambda-\alpha} = 0$, and our assertion is trivial. Assume that our assertion is true for $n+1$. Set

$$V_n^{\mathbf{A}} = (f_i^n V(\lambda))_{\lambda-\alpha}^{\mathbf{A}}, \quad L_n = f_i^n V(\lambda) \cap L(\lambda)_{\lambda-\alpha} \quad \text{and} \quad E_n = V_n^{\mathbf{A}} \cap L_n \cap L_n^-.$$

If $\langle h_i, \lambda - \alpha + n\alpha_i \rangle = 0$, then $f_i^n(V(\lambda)_{\lambda - \alpha + n\alpha_i}) = 0$, and our assertion is trivial. Hence we may assume that $\langle h_i, \lambda - \alpha + n\alpha_i \rangle > 0$, in which case, $\tilde{f}_i^n: B(\lambda)_{\lambda - \alpha + n\alpha_i} \rightarrow B(\lambda)_{\lambda - \alpha}$ is injective.

Since $n \geq 1$, Lemma 10.3(a) gives $V(\lambda)_{\lambda - \alpha + n\alpha_i}^{\mathbf{A}} = \bigoplus_{b \in B(\lambda)_{\lambda - \alpha + n\alpha_i}} \mathbf{A}G_\lambda(b)$ and hence we have

$$V_n = \sum_{b \in B(\lambda)_{\lambda - \alpha + n\alpha_i}} \mathbf{A}f_i^n G_\lambda(b).$$

If $b \in B(\lambda)_{\lambda - \alpha + n\alpha_i}$ satisfies $\tilde{e}_i b \neq 0$, then $\mathbf{Z}(r-1)$ implies that $G_\lambda(b) \in (f_i V(\lambda))^{\mathbf{A}} = f_i V(\lambda)^{\mathbf{A}}$, which yields

$$f_i^n G_\lambda(b) \in V_{n+1}^{\mathbf{A}} := (f_i^{n+1} V(\lambda))^{\mathbf{A}}_{\lambda - \alpha}.$$

Set $B_0 := \{b \in B(\lambda)_{\lambda - \alpha + n\alpha_i} ; \tilde{e}_i b = 0\}$ and $F_0 := \sum_{b \in B_0} \mathbf{Q}f_i^n G_\lambda(b)$. Then we get

$$(10.2) \quad V_n^{\mathbf{A}} = \mathbf{A}F_0 + V_{n+1}^{\mathbf{A}}.$$

Note that

$$f_i^n G_\lambda(b) \in E_n, \quad f_i^n G_\lambda(b) \equiv \tilde{f}_i^n b \pmod{qL(\lambda)} \quad \text{for every } b \in B_0.$$

Set $F = F_0 + E_{n+1}$. Then $F \subset E_n$, and $V_{n+1}^{\mathbf{A}} = \mathbf{A}E_{n+1}$ by the induction hypothesis. Hence (10.2) implies $V_n^{\mathbf{A}} = \mathbf{A}F$.

Consider the morphism

$$F \rightarrow L_n/qL_n \simeq \left(\bigoplus_{b \in B_0} \mathbf{Q}\tilde{f}_i^n b \right) \oplus (L_{n+1}/qL_{n+1}).$$

Since $F_0 \rightarrow \bigoplus_{b \in B_0} \mathbf{Q}\tilde{f}_i^n b$ and $E_{n+1} \rightarrow L_{n+1}/qL_{n+1}$ are isomorphisms, $F \rightarrow L_n/qL_n$ is an isomorphism. Similarly, $F \rightarrow (L_n)^-/q^{-1}(L_n)^-$ is an isomorphism. Hence Lemma 8.4 implies the desired result. $\square$

Corollary 10.6. *For every $i \in I$, $\alpha \in Q_+(r)$ and $n \geq 1$, we have the following canonical isomorphism:*

$$\begin{aligned} (f_i^n U_q^-(\mathfrak{g}))_{-\alpha}^{\mathbf{A}} \cap L(\infty) \cap L(\infty)^- &\xrightarrow{\sim} \frac{(f_i^n U_q^-(\mathfrak{g}))_{-\alpha}^{\mathbf{A}} \cap L(\infty)}{(f_i^n U_q^-(\mathfrak{g}))_{-\alpha}^{\mathbf{A}} \cap qL(\infty)} \\ &\simeq \bigoplus_{b \in B(\infty)_{-\alpha} \cap \tilde{f}_i^n B(\infty)} \mathbf{Q}b. \end{aligned}$$

Proof. It suffices to observe that, for $\lambda \gg 0$, we have

$$\begin{aligned} U_q^-(\mathfrak{g})_{-\alpha} &\xrightarrow{\sim} V(\lambda)_{\lambda-\alpha}, & (f_i^n U_q^-(\mathfrak{g}))_{-\alpha}^{\mathbf{A}} &\xrightarrow{\sim} (f_i^n V(\lambda))_{\lambda-\alpha}^{\mathbf{A}}, \\ L(\infty)_{-\alpha} &\xrightarrow{\sim} L(\lambda)_{\lambda-\alpha}, & L(\infty)_{-\alpha}^- &\xrightarrow{\sim} L(\lambda)_{\lambda-\alpha}^-, \\ \bigoplus_{b \in B(\infty)_{-\alpha} \cap \tilde{f}_i^n B(\infty)} \mathbf{Q}b &\xrightarrow{\sim} \bigoplus_{b \in B(\lambda)_{\lambda-\alpha} \cap \tilde{f}_i^n B(\lambda)} \mathbf{Q}b. \end{aligned}$$

□

For $\alpha \in Q_+(r)$, let

$$G_i: \bigoplus_{b \in B(\infty)_{-\alpha} \cap \tilde{f}_i B(\infty)} \mathbf{Q}b \longrightarrow (f_i U_q^-(\mathfrak{g}))_{-\alpha}^{\mathbf{A}} \cap L(\infty) \cap L(\infty)^-$$

be the inverse of the canonical isomorphism given in Corollary 10.6 (with $n = 1$).

Then we have

$$(f_i U_q^-(\mathfrak{g}))_{-\alpha}^{\mathbf{A}} = \bigoplus_{b \in B(\infty)_{-\alpha} \cap \tilde{f}_i B(\infty)} \mathbf{A}G_i(b).$$

Now we will prove:

Lemma 10.7. *For $i, j \in I$, $\alpha \in Q_+(r)$ and $b \in B(\infty)_{-\alpha} \cap \tilde{f}_i B(\infty) \cap \tilde{f}_j B(\infty)$, we have*

$$G_i(b) = G_j(b).$$

Proof. Let $b = \tilde{f}_i \cdots \tilde{f}_k \cdot 1 \in B(\infty)_{-\alpha} \cap \tilde{f}_i B(\infty) \cap \tilde{f}_j B(\infty)$. Take $\lambda \in P^+$ such that $\lambda(h_k) = 0$ and $\lambda(h_p) \gg 0$ for $p \in I \setminus \{k\}$. Since $f_k v_\lambda = 0$, we get $\bar{\pi}_\lambda(b) = 0$, which implies

$$G_i(b)v_\lambda \equiv 0 \pmod{qL(\lambda)}.$$

That is, $G_i(b)v_\lambda \in (f_i V(\lambda))_{\lambda-\alpha}^{\mathbf{A}} \cap qL(\lambda) \cap L(\lambda)^-$.

By Proposition 10.4, we must have $G_i(b)v_\lambda = 0$ in $V(\lambda)$. Note that

$$\begin{aligned} V(\lambda)_{\lambda-\alpha} &\simeq U_q^-(\mathfrak{g})_{-\alpha} / (U_q^-(\mathfrak{g})_{-\alpha+\alpha_k} f_k + \sum_{p \in I^{\text{re}} \setminus \{k\}} U_q^-(\mathfrak{g})_{-\alpha+(\lambda(h_p)+1)\alpha_i} f_p^{(\lambda(h_p)+1)}) \\ &\simeq U_q^-(\mathfrak{g})_{-\alpha} / U_q^-(\mathfrak{g})_{-\alpha+\alpha_k} f_k \end{aligned}$$

as a $U_q^-(\mathfrak{g})$ -module. It follows that $G_i(b) \in U_q^-(\mathfrak{g})f_k \cap U_{\mathbf{A}}^-(\mathfrak{g})$. By applying the anti-automorphism $\star$ on $G_i(b)$, we have

$$G_i(b)^\star \in f_k U_q^-(\mathfrak{g}) \cap U_{\mathbf{A}}^-(\mathfrak{g}) = (f_k U_q(\mathfrak{g}))^{\mathbf{A}}.$$

Moreover, by Corollary 7.40, we have $G_i(b)^\star \in (f_k U_q^-(\mathfrak{g}))_{-\alpha}^{\mathbf{A}} \cap L(\infty) \cap L(\infty)^-$.

Similarly, $G_j(b)$ satisfies $G_j(b)^\star \in (f_k U_q^-(\mathfrak{g}))^{\mathbf{A}} \cap L(\infty) \cap L(\infty)^-$. Since $G_i(b) \equiv G_j(b) \equiv b \pmod{qL(\infty)}$, we have $G_i(b)^\star \equiv G_j(b)^\star \pmod{qL(\infty)}$. Hence, Corollary 10.6 implies $G_i(b)^\star = G_j(b)^\star$, which yields $G_i(b) = G_j(b)$ as desired. $\square$

Let $b \in B(\infty)_{-\alpha}$ with $\alpha \in Q_+(r)$. Since $r \geq 1$, we have $\tilde{e}_i b \neq 0$ for some $i \in I$; that is, $b \in B(\infty)_{-\alpha} \cap \tilde{f}_i B(\infty)$ for some $i \in I$. By Lemma 10.7, we may define a map

$$G_\infty: L(\infty)_{-\alpha}/qL(\infty)_{-\alpha} \longrightarrow U_q^-(\mathfrak{g})_{-\alpha}^{\mathbf{A}} \cap L(\infty) \cap L(\infty)^-$$

by $G_\infty(b) = G_i(b)$ for any i with $\tilde{e}_i b \neq 0$. By the definition, we have

$$(10.3) \quad \begin{aligned} G_\infty(b) &\equiv b \pmod{qL(\infty)}, \\ (f_i^n U_q^-(\mathfrak{g}))_{-\alpha}^{\mathbf{A}} &= \bigoplus_{b \in B(\infty)_{-\alpha} \cap \tilde{f}_i^n B(\infty)} \mathbf{A} G_\infty(b) \quad \text{for } n \geq 1. \end{aligned}$$

Set $F = \sum_{b \in B(\infty)_{-\alpha}} \mathbf{Q} G_\infty(b) \subset U_q^-(\mathfrak{g})_{-\alpha}^{\mathbf{A}} \cap L(\infty) \cap L(\infty)^-$. Since $U_{\mathbf{A}}^-(\mathfrak{g})_{-\alpha} = \sum_i (f_i U_q^-(\mathfrak{g}))_{-\alpha}^{\mathbf{A}}$, we obtain

$$U_{\mathbf{A}}^-(\mathfrak{g})_{-\alpha} = \mathbf{A}F.$$

Moreover, the canonical maps

$$F \rightarrow L(\infty)_{-\alpha}/qL(\infty)_{-\alpha} \text{ and } F \rightarrow L(\infty)_{-\alpha}^-/q^{-1}L(\infty)_{-\alpha}^-$$

are bijective. Therefore, by Lemma 8.4, we conclude $\mathbf{X}(r)$.

Let us prove $\mathbf{Y}(r)$. Note that for all $\alpha \in Q_+(r)$, $b \in B(\infty)_{-\alpha}$ and $\lambda \in P^+$, if $\pi_\lambda(b) = 0$, then $G_\infty(b)v_\lambda = 0$. Indeed, let $i \in I$ be such that $\tilde{e}_i b \neq 0$. Then we have

$$G_\infty(b)v_\lambda \in (f_i V(\lambda))_{\lambda-\alpha}^{\mathbf{A}} \cap qL(\lambda) \cap L(\lambda)^-,$$

and by Proposition 10.4, we obtain $G_\infty(b)v_\lambda = 0$.

Set $F := \sum_{\substack{b \in B(\infty)_{-\alpha} \\ \pi_\lambda(b) \neq 0}} \mathbf{Q} G_\infty(b)v_\lambda \subset V(\lambda)_{\lambda-\alpha}^{\mathbf{A}} \cap L(\lambda) \cap L(\lambda)^-$. Then we have $V(\lambda)_{\lambda-\alpha}^{\mathbf{A}} = \mathbf{A}F$. By Proposition 7.34, the homomorphisms $F \rightarrow L(\lambda)_{\lambda-\alpha}/qL(\lambda)_{\lambda-\alpha}$ and $F \rightarrow$

$L(\lambda)_{\lambda-\alpha}^-/q^{-1}L(\lambda)_{\lambda-\alpha}^-$ are isomorphisms. Therefore, by Lemma 8.4, we obtain the statement $\mathbf{Y}(r)$.

Finally, the statement $\mathbf{Z}(r)$ follows from Proposition 10.4, which completes the proof.

Remark 10.8.

- (i) We have $G_\infty(b)v_\lambda = G_\lambda(\pi_\lambda(b))$ for any $\lambda \in P^+$ and $b \in B(\infty)$.
- (ii) If $i \in I^{\text{im}}$, we have

$$G_\lambda(\tilde{f}_i^n b) = f_i^n G_\lambda(b) \text{ for every } b \in B(\lambda) \text{ and } n \in \mathbf{Z}_{\geq 0}.$$

- (iii) We have

$$(f_i^n V(\lambda))^{\mathbf{A}} = f_i^n V(\lambda) \cap V(\lambda)^{\mathbf{A}}.$$

Indeed, $\{G_\lambda(b)\}_{b \in \tilde{f}_i^n B(\lambda)}$ is an $\mathbf{A}$ -basis of $(f_i^n V(\lambda))^{\mathbf{A}}$, and hence it is also a $\mathbf{Q}(q)$ -basis of $f_i^n V(\lambda)$.

REFERENCES

- [1] G. Benkart, S.-J. Kang, D. J. Melville, *Quantized enveloping algebras for Borchersd's superalgebras*, Trans. Amer. Math. Soc. **350** (1998), 3297–3319.
- [2] R. E. Borcherds, *Generalized Kac-Moody algebras*, J. Algebra **115** (1988), 501–512.
- [3] ———, *Monstrous moonshine and monstrous Lie superalgebras*, Invent. Math. **109** (1992), 405–444.
- [4] C. W. Curtis, I. Reiner, *Methods of Representation Theory, Vol. I*, John Wiley & Sons Inc., New York, 1981.
- [5] V. G. Drinfel'd, *Hopf algebra and the Yang-Baxter equation*, Soviet Math. Dokl. **32** (1985), 254–258.
- [6] J. Hong, S.-J. Kang, *Introduction to Quantum Groups and Crystal Bases*, Graduate Studies in Mathematics **42**, Amer. Math. Soc., 2002.
- [7] M. Jimbo, *A q -difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation*, Lett. Math. Phys. **10** (1985), 63–69.
- [8] S.-J. Kang, *Quantum deformations of generalized Kac-Moody algebras and their modules*, J. Algebra **175** (1995), 1041–1066.
- [9] M. Kashiwara, *Crystallizing the q -analogue of universal enveloping algebras*, Comm. Math. Phys. **133** (1990), 249–260.
- [10] ———, *On crystal bases of the q -analogue of universal enveloping algebras*, Duke Math. J. **63** (1991), 465–516.

- [11] ———, *Bases cristallines des groupes quantiques*, (rédigé par Charles Cochet), Cours spécialisés **9**, Société Mathématique de France, 2002.
- [12] G. Lusztig, *Quantum deformation of certain simple modules over enveloping algebras*, Adv. Math. **70** (1988), 237–249.
- [13] ———, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), 447–498.

◊DEPARTMENT OF MATHEMATICS, NS30, SEOUL NATIONAL UNIVERSITY, SEOUL 151-747, KOREA

E-mail address: khjeong@math.snu.ac.kr

*SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, 207-43 CHEONGRYANGRI-DONG, DONGDAEMUN-GU, SEOUL 130-722, KOREA

E-mail address: sjkang@kias.re.kr

†RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO 606, JAPAN

E-mail address: masaki@kurims.kyoto-u.ac.jp